wwPDB X-ray Structure Validation Summary Report (i)

Aug 4, 2026 – 08:15 PM EDT

PDB ID	:	3ABQ / pdb_00003abq
Title	:	Crystal structure of ethanolamine ammonia-lyase from Escherichia coli complexed with CN-Cbl and 2-amino-1-propanol
Authors	:	Shibata, N.
Deposited on	:	2009-12-21
Resolution	:	2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	2%, 93%, 6%
1	C	453	3%, 89%, 10%, •
2	B	306	14%, 70%, 10%, •, 18%
2	D	306	43%, 62%, 17%, •, 19%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 12057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ethanolamine ammonia-lyase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	453	3500	2197	599	681	23	0	7	0
1	C	453	3497	2194	599	680	24	8	6	0

- Molecule 2 is a protein called Ethanolamine ammonia-lyase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	252	1916	1197	347	362	10	0	0	0
2	D	248	1891	1183	342	357	9	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	MET	-	expression tag	UNP P19636
B	-9	ASP	-	expression tag	UNP P19636
B	-8	GLN	-	expression tag	UNP P19636
B	-7	SER	-	expression tag	UNP P19636
B	-6	SER	-	expression tag	UNP P19636
B	-5	HIS	-	expression tag	UNP P19636
B	-4	HIS	-	expression tag	UNP P19636
B	-3	HIS	-	expression tag	UNP P19636
B	-2	HIS	-	expression tag	UNP P19636
B	-1	HIS	-	expression tag	UNP P19636
B	0	HIS	-	expression tag	UNP P19636
D	-10	MET	-	expression tag	UNP P19636
D	-9	ASP	-	expression tag	UNP P19636
D	-8	GLN	-	expression tag	UNP P19636
D	-7	SER	-	expression tag	UNP P19636
D	-6	SER	-	expression tag	UNP P19636

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	expression tag	UNP P19636
D	-4	HIS	-	expression tag	UNP P19636
D	-3	HIS	-	expression tag	UNP P19636
D	-2	HIS	-	expression tag	UNP P19636
D	-1	HIS	-	expression tag	UNP P19636
D	0	HIS	-	expression tag	UNP P19636

- Molecule 3 is (2S)-2-aminopropan-1-ol (CCD ID: 2A1) (formula: C_3H_9NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	5	3	1	1	0	0
			Total	C	N	O		
3	C	1	5	3	1	1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
4	A	1	6	3	3	0	0
4	A	1	6	3	3	0	0
4	A	1	6	3	3	0	0
4	C	1	6	3	3	0	0
4	C	1	6	3	3	0	0
4	C	1	6	3	3	0	0
4	C	1	6	3	3	0	0

- Molecule 5 is COBALAMIN (CCD ID: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Co	N	O	P		
5	B	1	91	62	1	13	14	1	0	0
			Total	C	Co	N	O	P		
5	D	1	94	63	1	14	15	1	0	1

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	422	422	422	0	0
			Total	O		
6	B	124	124	124	0	0
			Total	O		
6	C	377	377	377	0	0
			Total	O		
6	D	87	87	87	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ethanolamine ammonia-lyase heavy chain

Chain A:





• Molecule 1: Ethanolamine ammonia-lyase heavy chain

Chain C:



• Molecule 2: Ethanolamine ammonia-lyase light chain

Chain B:



• Molecule 2: Ethanolamine ammonia-lyase light chain

Chain D:

43%
62%
17%
19%


MET ASP GLN SER SER HIS HIS HIS HIS HIS HIS MET ASP GLN LYS GLN ILE GLU GLU ILE VAL ARG SER VAL MET ALA SER MET GLY GLN ALA ALA PRO ALA PRO SER GLU ALA LYS CYS ALA THR THR ASN CYS ALA ALA PRO VAL THR SER GLU SER CYS A44 W55 D65 V66
E69 L70 R71 R72 S73 R77 H99 K103 D104 T105 V106 L107 K108 E109 V110 P111 E112 E113 W114 V115 K116 A117 Q118 G119 L120 L121 E122 V123 R124 S125 E126 I127 S128 Y133 L134 T135 R136 R141 R142 L143 C144 ALA E146 A147 V148 E149 A150 L151 K152 ALA GLN CYS V156 A157
N158 P159 D160 V161 Q162 V163 V164 I165 S170 T171 D172 A173 I174 T175 Y178 E179 E180 I181 L182 P183 P184 L185 M186 A187 G188 L189 K190 Q191 A192 G193 L194 K195 V196 G197 T198 P199 F200 F201 V202 R203 Y204 G205 R206 V207 K208 I209 I213 G214 E215 I216 L217 G218 A219 K220 V221 V222 I223
L224 L225 E228 R229 L232 G233 Q234 L238 S239 C240 Y241 A242 V243 Y244 S245 P246 R247 M248 A249 T250 T251 V252 T257 C258 I259 I262 H263 Q264 G265 G266 T267 P268 P269 V270 E271 A272 A273 A274 V275 I276 V277 D278 L279 A280 K281 R282 M283 L284 E285 Q286 K287 A288 S289 G290 I291
N292 M293 T294 R295

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α, β, γ	241.87Å 241.87Å 76.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.05 50.00 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-2.05) 99.4 (50.00-2.05)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.65 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.5.0087	Depositor
R, R_{free}	0.230 , 0.269 0.255 , 0.290	Depositor DCC
R_{free} test set	7958 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	24.2	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 40.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.42$, $< L^2 > = 0.25$	Xtriage
Estimated twinning fraction	0.049 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.89	EDS
Total number of atoms	12057	wwPDB-VP
Average B, all atoms (Å^2)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.15% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, B12, 2A1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.31	0/3575	0.66	2/4839 (0.0%)
1	C	0.31	0/3569	0.66	2/4830 (0.0%)
2	B	0.30	0/1943	0.68	0/2633
2	D	0.30	0/1916	0.67	0/2593
All	All	0.31	0/11003	0.67	4/14895 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	331	GLY	CA-C-N	5.62	125.24	119.56
1	A	331	GLY	C-N-CA	5.62	125.24	119.56
1	C	331	GLY	CA-C-N	5.26	124.87	119.56
1	C	331	GLY	C-N-CA	5.26	124.87	119.56

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	326	VAL	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3500	0	3484	19	0
1	C	3497	0	3477	42	0
2	B	1916	0	1973	19	0
2	D	1891	0	1948	27	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
4	A	24	0	32	0	0
4	C	24	0	32	5	0
5	B	91	0	88	16	0
5	D	94	0	8	3	0
6	A	422	0	0	2	0
6	B	124	0	0	0	0
6	C	377	0	0	2	0
6	D	87	0	0	1	0
All	All	12057	0	11042	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:195[B]:VAL:HG12	1:C:401[B]:MET:CE	1.37	1.51
1:C:195[B]:VAL:CG1	1:C:401[B]:MET:CE	2.11	1.28
1:C:195[B]:VAL:CG1	1:C:401[B]:MET:HE3	1.68	1.19
1:C:195[B]:VAL:HG12	1:C:401[B]:MET:HE1	1.20	1.09
5:B:601:B12:H362	5:B:601:B12:H351	1.34	1.05

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/453 (101%)	443 (97%)	15 (3%)	0	100	100
1	C	457/453 (101%)	444 (97%)	13 (3%)	0	100	100
2	B	250/306 (82%)	245 (98%)	5 (2%)	0	100	100
2	D	242/306 (79%)	236 (98%)	6 (2%)	0	100	100
All	All	1407/1518 (93%)	1368 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/370 (102%)	372 (99%)	5 (1%)	61	63
1	C	376/370 (102%)	367 (98%)	9 (2%)	43	40
2	B	206/251 (82%)	193 (94%)	13 (6%)	16	9
2	D	204/251 (81%)	186 (91%)	18 (9%)	9	4
All	All	1163/1242 (94%)	1118 (96%)	45 (4%)	29	23

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	113	GLU
2	D	223	ILE
2	D	123	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	182	LEU
2	D	232	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	349	HIS
2	D	99	HIS
2	D	177	ASN
2	B	131	ASN
2	B	177	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	B12	D	601[B]	-	94,101,101	1.42	13 (13%)	149,166,166	1.57	29 (19%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	C	2008	-	5,5,5	0.36	0	5,5,5	0.28	0
3	2A1	A	602	-	4,4,4	0.64	0	1,4,4	0.28	0
4	GOL	A	2006	-	5,5,5	0.38	0	5,5,5	0.32	0
4	GOL	C	2003	-	5,5,5	0.35	0	5,5,5	0.41	0
5	B12	B	601	-	94,101,101	1.50	13 (13%)	149,166,166	1.41	25 (16%)
5	B12	D	601[A]	-	94,101,101	1.42	13 (13%)	149,166,166	1.57	29 (19%)
3	2A1	C	602	-	4,4,4	0.65	0	1,4,4	0.16	0
4	GOL	A	2007	-	5,5,5	0.36	0	5,5,5	0.35	0
4	GOL	C	2005	-	5,5,5	0.37	0	5,5,5	0.29	0
4	GOL	C	2004	-	5,5,5	0.37	0	5,5,5	0.31	0
4	GOL	A	2002	-	5,5,5	0.41	0	5,5,5	0.40	0
4	GOL	A	2001	-	5,5,5	0.37	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	B12	D	601[B]	-	-	20/56/223/223	0/3/11/11
4	GOL	C	2008	-	-	2/4/4/4	-
3	2A1	A	602	-	-	1/2/2/2	-
4	GOL	A	2006	-	-	2/4/4/4	-
4	GOL	C	2003	-	-	3/4/4/4	-
5	B12	B	601	-	-	20/56/223/223	0/3/11/11
5	B12	D	601[A]	-	-	19/56/223/223	0/3/11/11
3	2A1	C	602	-	-	0/2/2/2	-
4	GOL	A	2007	-	-	0/4/4/4	-
4	GOL	C	2005	-	-	2/4/4/4	-
4	GOL	C	2004	-	-	2/4/4/4	-
4	GOL	A	2002	-	-	2/4/4/4	-
4	GOL	A	2001	-	-	1/4/4/4	-

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	601	B12	C19-N24	-6.10	1.41	1.49
5	D	601[A]	B12	C19-N24	-5.92	1.41	1.49
5	D	601[B]	B12	C19-N24	-5.92	1.41	1.49
5	B	601	B12	C8B-C9B	5.01	1.48	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	601[A]	B12	C8B-C9B	4.98	1.48	1.40

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	D	601[A]	B12	C47-C12-C46	6.87	120.79	109.41
5	D	601[B]	B12	C47-C12-C46	6.87	120.79	109.41
5	B	601	B12	C19-N24-C16	4.42	112.12	107.29
5	D	601[A]	B12	C54-C17-C18	-4.41	106.66	112.99
5	D	601[B]	B12	C54-C17-C18	-4.41	106.66	112.99

There are no chirality outliers.

5 of 74 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2002	GOL	O1-C1-C2-O2
4	A	2002	GOL	O1-C1-C2-C3
4	C	2004	GOL	C1-C2-C3-O3
4	C	2005	GOL	O1-C1-C2-O2
4	C	2005	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	601[B]	B12	1	0
4	C	2008	GOL	3	0
5	B	601	B12	16	0
5	D	601[A]	B12	2	0
4	C	2005	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.

Ligand B12 D 601 (B)



Ligand B12 B 601



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	453/453 (100%)	0.04	7 (1%)	72	73	15, 27, 46, 63	7 (1%)
1	C	453/453 (100%)	0.19	14 (3%)	51	52	11, 29, 52, 75	6 (1%)
2	B	252/306 (82%)	1.21	43 (17%)	4	3	21, 53, 77, 100	0
2	D	248/306 (81%)	2.29	131 (52%)	0	0	18, 75, 120, 146	0
All	All	1406/1518 (92%)	0.70	195 (13%)	6	6	11, 35, 95, 146	13 (0%)

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	192	ALA	11.5
2	D	194	LEU	7.8
2	D	156	VAL	7.5
2	B	192	ALA	6.6
2	D	44	ALA	6.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	C	2008	6/6	0.83	0.15	37,44,47,57	0
4	GOL	C	2004	6/6	0.85	0.14	45,46,52,54	0
4	GOL	A	2002	6/6	0.87	0.15	27,36,40,43	0
4	GOL	A	2006	6/6	0.90	0.16	27,40,46,46	0
4	GOL	C	2003	6/6	0.90	0.15	30,37,44,48	0
4	GOL	C	2005	6/6	0.91	0.13	31,39,47,60	0
3	2A1	C	602	5/5	0.91	0.17	23,23,34,41	0
4	GOL	A	2001	6/6	0.92	0.11	25,28,32,39	0
3	2A1	A	602	5/5	0.93	0.13	22,22,37,41	0
5	B12	D	601[A]	91/91	0.93	0.13	14,37,51,57	3
5	B12	D	601[B]	91/91	0.93	0.13	14,37,51,57	3
5	B12	B	601	91/91	0.95	0.10	18,28,40,49	0
4	GOL	A	2007	6/6	0.97	0.09	19,24,24,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B12 D 601 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B12 D 601 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B12 B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.