



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 25, 2026 – 09:16 am BST

PDB ID : 4AYB / pdb_00004ayb
Title : RNAP at 3.2Å
Authors : Wojtas, M.N.; Mogni, M.; Millet, O.; Bell, S.D.; Abrescia, N.G.A.
Deposited on : 2012-06-19
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	880	
2	B	1131	
3	C	395	
4	D	265	
5	E	180	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	113	
7	G	132	
8	H	84	
9	K	95	
10	L	92	
11	N	66	
12	P	48	
13	Q	104	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
16	SF4	D	1264	-	-	X	-

2 Entry composition [i](#)

There are 16 unique types of molecules in this entry. The entry contains 54890 atoms, of which 27769 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	872	Total	C	H	N	O	S	0	0	0
			13986	4424	7029	1225	1282	26			

- Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	1103	Total	C	H	N	O	S	0	0	0
			17664	5548	8908	1552	1627	29			

- Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	376	Total	C	H	N	O	S	0	0	0
			5974	1840	3068	493	564	9			

- Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	262	Total	C	H	N	O	S	0	0	0
			4215	1339	2128	337	398	13			

- Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
5	E	171	Total	C	H	N	O	S	0	0	0
			2772	874	1413	229	251	5			

- Molecule 6 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
6	F	105	Total	C	H	N	O	S	0	0	0
			1666	519	839	134	171	3			

- Molecule 7 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
7	G	113	Total	C	H	N	O	S	0	0	0
			1816	572	915	152	173	4			

- Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
8	H	76	Total	C	H	N	O		0	0	0
			1284	405	660	111	108				

- Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
9	K	84	Total	C	H	N	O	S	0	0	0
			1390	431	717	123	118	1			

- Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
10	L	91	Total	C	H	N	O	S	0	0	0
			1449	454	742	114	137	2			

- Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
11	N	65	Total	C	H	N	O	S	0	0	0
			1058	332	537	94	88	7			

- Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
12	P	44	Total	C	H	N	O	S	0	0	0
			744	236	387	62	54	5			

- Molecule 13 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
13	Q	50	Total	C	H	N	O	S	0	0	0
			854	270	426	74	83	1			


- Molecule 14 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	A	4	Total Zn 4 4	0	0
14	B	3	Total Zn 3 3	0	0
14	C	1	Total Zn 1 1	0	0
14	N	1	Total Zn 1 1	0	0
14	P	1	Total Zn 1 1	0	0

- Molecule 15 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	A	1	Total Mg 1 1	0	0

- Molecule 16 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	D	1	Total Fe S 7 3 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA-DIRECTED RNA POLYMERASE









• Molecule 3: DNA-DIRECTED RNA POLYMERASE




• Molecule 4: DNA-DIRECTED RNA POLYMERASE




• Molecule 5: DNA-DIRECTED RNA POLYMERASE







• Molecule 6: DNA-DIRECTED RNA POLYMERASE




• Molecule 7: DNA-DIRECTED RNA POLYMERASE



• Molecule 8: DNA-DIRECTED RNA POLYMERASE



• Molecule 9: DNA-DIRECTED RNA POLYMERASE





• Molecule 10: DNA-DIRECTED RNA POLYMERASE





- Molecule 11: DNA-DIRECTED RNA POLYMERASE



- Molecule 12: DNA-DIRECTED RNA POLYMERASE



- Molecule 13: DNA-DIRECTED RNA POLYMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	195.72Å 212.41Å 128.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.10 – 3.20 40.10 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (40.10-3.20) 99.4 (40.10-3.20)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 3.18Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.242 , 0.300 0.236 , 0.294	Depositor DCC
R_{free} test set	4428 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	70.6	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 80.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	54890	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SF4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/7108	0.72	1/9618 (0.0%)
2	B	0.32	0/8923	0.70	0/12071
3	C	0.32	0/2930	0.75	0/3944
4	D	0.28	0/2123	0.66	0/2870
5	E	0.27	0/1379	0.66	0/1861
6	F	0.32	0/836	0.70	0/1133
7	G	0.28	0/913	0.72	0/1224
8	H	0.29	0/638	0.70	0/864
9	K	0.33	0/682	0.72	1/921 (0.1%)
10	L	0.27	0/717	0.65	0/968
11	N	0.32	0/532	0.71	0/718
12	P	0.36	0/365	0.67	0/489
13	Q	0.28	0/434	0.63	1/580 (0.2%)
All	All	0.32	0/27580	0.70	3/37261 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	811	VAL	N-CA-C	6.34	116.45	110.30
9	K	85	GLY	N-CA-C	-6.20	107.70	115.08
13	Q	56	ASN	N-CA-C	5.06	116.88	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6957	7029	7013	429	0
2	B	8756	8908	8888	454	0
3	C	2906	3068	3063	208	0
4	D	2087	2128	2125	68	0
5	E	1359	1413	1411	80	0
6	F	827	839	839	88	0
7	G	901	915	912	88	0
8	H	624	660	658	26	0
9	K	673	717	716	23	0
10	L	707	742	739	20	0
11	N	521	537	535	42	0
12	P	357	387	387	26	0
13	Q	428	426	426	40	0
14	A	4	0	0	0	0
14	B	3	0	0	0	0
14	C	1	0	0	0	0
14	N	1	0	0	0	0
14	P	1	0	0	0	0
15	A	1	0	0	0	0
16	D	7	0	0	4	0
All	All	27121	27769	27712	1481	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 1481 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:93:ARG:CB	6:F:94:THR:HA	1.87	1.05
1:A:146:CYS:SG	1:A:154:PHE:CZ	2.55	1.00
2:B:221:PRO:HB2	2:B:222:GLY:HA2	1.41	1.00
6:F:93:ARG:HB3	6:F:94:THR:HA	1.47	0.95
1:A:683:GLU:HA	1:A:684:LEU:HB3	1.51	0.93

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	868/880 (99%)	712 (82%)	124 (14%)	32 (4%)	2	18
2	B	1097/1131 (97%)	928 (85%)	133 (12%)	36 (3%)	3	21
3	C	372/395 (94%)	304 (82%)	49 (13%)	19 (5%)	1	13
4	D	260/265 (98%)	218 (84%)	37 (14%)	5 (2%)	6	33
5	E	167/180 (93%)	150 (90%)	16 (10%)	1 (1%)	21	56
6	F	103/113 (91%)	78 (76%)	17 (16%)	8 (8%)	1	5
7	G	111/132 (84%)	83 (75%)	20 (18%)	8 (7%)	1	6
8	H	74/84 (88%)	63 (85%)	8 (11%)	3 (4%)	2	17
9	K	82/95 (86%)	69 (84%)	9 (11%)	4 (5%)	1	13
10	L	89/92 (97%)	83 (93%)	4 (4%)	2 (2%)	5	29
11	N	63/66 (96%)	47 (75%)	10 (16%)	6 (10%)	0	3
12	P	42/48 (88%)	29 (69%)	9 (21%)	4 (10%)	0	3
13	Q	48/104 (46%)	38 (79%)	5 (10%)	5 (10%)	0	2
All	All	3376/3585 (94%)	2802 (83%)	441 (13%)	133 (4%)	2	18

5 of 133 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	ASP
1	A	43	SER
1	A	44	VAL
1	A	229	ILE
1	A	291	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	758/766 (99%)	721 (95%)	37 (5%)	22	55
2	B	951/975 (98%)	903 (95%)	48 (5%)	22	55
3	C	324/341 (95%)	305 (94%)	19 (6%)	18	50
4	D	235/238 (99%)	234 (100%)	1 (0%)	84	86
5	E	150/158 (95%)	142 (95%)	8 (5%)	20	53
6	F	99/107 (92%)	89 (90%)	10 (10%)	7	30
7	G	106/125 (85%)	99 (93%)	7 (7%)	15	47
8	H	69/75 (92%)	68 (99%)	1 (1%)	59	77
9	K	74/83 (89%)	73 (99%)	1 (1%)	59	77
10	L	79/80 (99%)	79 (100%)	0	100	100
11	N	59/60 (98%)	58 (98%)	1 (2%)	53	75
12	P	40/43 (93%)	39 (98%)	1 (2%)	42	69
13	Q	48/96 (50%)	45 (94%)	3 (6%)	16	48
All	All	2992/3147 (95%)	2855 (95%)	137 (5%)	24	58

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	F	47	CYS
6	F	87	LEU
8	H	48	SER
2	B	230	MET
2	B	224	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	997	HIS
7	G	25	ASN
10	L	82	HIS
7	G	88	ASN
2	B	640	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 11 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
16	SF4	D	1264	4	0,9,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	SF4	D	1264	4	-	-	0/3/3/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	D	1264	SF4	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	872/880 (99%)	0.40	61 (6%)	22 14	50, 78, 123, 165	0
2	B	1103/1131 (97%)	0.40	53 (4%)	35 22	49, 77, 118, 173	0
3	C	376/395 (95%)	0.53	31 (8%)	17 11	52, 86, 126, 185	0
4	D	262/265 (98%)	0.38	10 (3%)	44 27	67, 87, 118, 140	0
5	E	171/180 (95%)	1.10	26 (15%)	5 4	63, 123, 158, 178	0
6	F	105/113 (92%)	1.24	14 (13%)	7 6	94, 137, 169, 215	0
7	G	113/132 (85%)	1.10	16 (14%)	6 5	69, 103, 127, 157	0
8	H	76/84 (90%)	0.46	5 (6%)	24 15	62, 79, 104, 121	0
9	K	84/95 (88%)	0.23	5 (5%)	27 17	58, 71, 111, 137	0
10	L	91/92 (98%)	0.18	1 (1%)	78 61	62, 84, 107, 159	0
11	N	65/66 (98%)	0.40	6 (9%)	14 9	62, 81, 102, 112	0
12	P	44/48 (91%)	1.21	11 (25%)	2 2	66, 88, 119, 152	0
13	Q	50/104 (48%)	0.70	3 (6%)	27 17	89, 115, 145, 158	0
All	All	3412/3585 (95%)	0.50	242 (7%)	22 14	49, 83, 135, 215	0

The worst 5 of 242 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	340	SER	6.2
5	E	72	PHE	4.9
7	G	92	TYR	4.9
1	A	122	LYS	4.9
2	B	476	MET	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
15	MG	A	1884	1/1	0.69	0.28	81,81,81,81	0
14	ZN	A	1882	1/1	0.90	0.25	81,81,81,81	0
14	ZN	B	2127	1/1	0.91	0.19	81,81,81,81	0
14	ZN	B	2128	1/1	0.93	0.25	81,81,81,81	0
14	ZN	A	1881	1/1	0.93	0.13	81,81,81,81	0
14	ZN	A	1883	1/1	0.94	0.15	81,81,81,81	0
14	ZN	P	1049	1/1	0.95	0.06	118,118,118,118	0
14	ZN	C	1395	1/1	0.95	0.07	81,81,81,81	0
16	SF4	D	1264	7/8	0.95	0.13	81,81,81,81	0
14	ZN	A	1880	1/1	0.96	0.05	81,81,81,81	0
14	ZN	B	2126	1/1	0.96	0.05	81,81,81,81	0
14	ZN	N	1066	1/1	0.97	0.06	88,88,88,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.