



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 3, 2026 – 09:54 PM EDT

PDB ID : 3BEN / pdb_00003ben
Title : Structure of N-(12-imidazolyl-dodecanoyl)-L-leucine inhibitor bound to the heme domain of Cytochrome P450-BM3
Authors : Tomchick, D.R.
Deposited on : 2007-11-19
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	<div> <div>4%</div> <div>90%</div> <div>7%</div> <div>.</div> </div>
1	B	470	<div> <div>8%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 8685 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 102.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	454	Total	C	N	O	S	0	15	0
			3778	2409	639	711	19			
1	B	457	Total	C	N	O	S	0	10	0
			3764	2402	640	702	20			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			27	21	3	3		
3	B	1	Total	C	N	O	0	0
			27	21	3	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	456	Total	O	0	0
			456	456		
6	B	448	Total	O	0	0
			448	448		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome P450 102



• Molecule 1: Cytochrome P450 102



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.82Å 148.21Å 63.79Å 90.00° 98.32° 90.00°	Depositor
Resolution (Å)	30.00 – 1.65 30.00 – 1.65	Depositor EDS
% Data completeness (in resolution range)	97.2 (30.00-1.65) 97.2 (30.00-1.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.68 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.161 , 0.191 0.162 , 0.191	Depositor DCC
R_{free} test set	3825 reflections (2.96%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8685	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, LEH, MG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/3863	0.89	1/5219 (0.0%)
1	B	0.86	2/3849 (0.1%)	0.93	5/5199 (0.1%)
All	All	0.82	2/7712 (0.0%)	0.91	6/10418 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	357	ILE	CA-CB	6.00	1.57	1.53
1	B	198	TYR	CG-CD2	5.71	1.51	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	230	SER	N-CA-C	-9.12	98.88	110.19
1	B	189	GLN	CA-C-N	-5.90	114.67	122.99
1	B	189	GLN	C-N-CA	-5.90	114.67	122.99
1	B	380	GLU	N-CA-C	5.55	118.10	111.71
1	A	230	SER	N-CA-C	-5.49	101.20	109.60
1	B	378	ARG	CB-CA-C	-5.38	100.02	110.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3778	0	3727	26	0
1	B	3764	0	3734	39	0
2	A	86	0	60	3	0
2	B	86	0	60	6	0
3	A	27	0	36	7	0
3	B	27	0	36	5	0
4	B	1	0	0	0	0
5	B	12	0	12	0	0
6	A	456	0	0	2	0
6	B	448	0	0	7	0
All	All	8685	0	7665	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:413[B]:VAL:HG13	6:B:690:HOH:O	1.48	1.12
1:A:78:VAL:HG21	3:A:500:LEH:H13	1.55	0.88
1:A:49[B]:THR:HG21	1:A:354[B]:MET:HG2	1.63	0.79
1:B:306:LYS:N	1:B:306:LYS:HD3	2.01	0.76
1:B:126[B]:LEU:HD11	1:B:145:MET:HE1	1.69	0.75
1:A:49[B]:THR:CG2	1:A:354[B]:MET:HG2	2.21	0.71
1:B:181:LEU:O	1:B:185[B]:MET:HG3	1.91	0.70
1:A:109:GLN:HB2	1:A:404[A]:GLN:OE1	1.91	0.69
1:A:185:MET:SD	3:A:500:LEH:H9A	2.33	0.68
1:B:118[A]:MET:HE1	6:B:880:HOH:O	1.94	0.67
1:B:413[B]:VAL:CG1	6:B:690:HOH:O	2.23	0.64
2:B:471[A]:HEM:HBC2	2:B:471[A]:HEM:HMC1	1.78	0.64
2:A:470[A]:HEM:HMC1	2:A:470[A]:HEM:HBC2	1.81	0.62
1:A:75:LEU:HD21	1:A:87:PHE:HE2	1.66	0.61
1:A:118[A]:MET:SD	6:A:711:HOH:O	2.56	0.59
1:B:313:TYR:HA	1:B:316:MET:HE3	1.85	0.57
1:A:437:LEU:HD23	1:A:437:LEU:H	1.69	0.57
1:B:75:LEU:HD21	1:B:87:PHE:HE1	1.70	0.57
1:B:306:LYS:N	1:B:306:LYS:CD	2.67	0.57
3:B:500:LEH:O3	3:B:500:LEH:H4	2.05	0.56
1:B:437:LEU:HB2	3:B:500:LEH:H12A	1.88	0.56
1:B:437:LEU:H	1:B:437:LEU:HD23	1.71	0.55
1:B:304:SER:HB2	1:B:306:LYS:HZ1	1.71	0.54
1:A:126[A]:LEU:HD21	1:A:145:MET:HE1	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181:LEU:O	1:B:185[A]:MET:HG2	2.09	0.53
1:B:409:GLU:O	1:B:413[B]:VAL:HG22	2.09	0.53
1:B:78:VAL:HG21	3:B:500:LEH:H13	1.90	0.52
1:B:118[A]:MET:HE3	1:B:156:CYS:HA	1.92	0.51
1:B:306:LYS:CD	1:B:306:LYS:H	2.24	0.51
1:B:229:GLN:C	1:B:230:SER:O	2.49	0.51
1:A:148:LEU:HD21	1:A:413[A]:VAL:HG21	1.92	0.51
1:A:26:VAL:CG2	3:A:500:LEH:H5	2.40	0.50
1:A:229:GLN:C	1:A:230:SER:O	2.51	0.50
1:B:304:SER:HB2	1:B:306:LYS:NZ	2.26	0.50
1:A:223:ARG:HD3	1:A:228:GLU:HG2	1.94	0.49
1:A:143:GLU:HG2	6:A:864:HOH:O	2.12	0.49
1:A:79:ARG:HG3	1:A:83:GLY:O	2.13	0.48
1:B:306:LYS:HD3	1:B:306:LYS:H	1.74	0.48
1:B:118[A]:MET:CE	1:B:156:CYS:HA	2.44	0.47
1:B:400:CYS:HA	2:B:471[A]:HEM:CHA	2.45	0.47
1:A:49[B]:THR:HG21	1:A:354[B]:MET:CG	2.39	0.47
1:A:185:MET:SD	3:A:500:LEH:C9	3.02	0.47
1:B:185[B]:MET:HE1	1:B:437:LEU:HA	1.97	0.47
1:B:367:TRP:HB2	1:B:371:VAL:HG12	1.97	0.46
1:B:128[A]:GLN:NE2	6:B:709:HOH:O	2.48	0.46
1:A:26:VAL:HG22	3:A:500:LEH:H5	1.98	0.45
1:A:183:GLU:HG3	1:A:187:LYS:HE3	1.99	0.45
1:B:269:THR:HA	2:B:471[A]:HEM:HAB	1.98	0.45
1:B:402:GLY:HA3	2:B:471[B]:HEM:C2B	2.51	0.45
1:B:158:PHE:CE1	1:B:258:ILE:HG12	2.53	0.44
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.99	0.44
1:A:75:LEU:HD23	3:A:500:LEH:H13A	1.99	0.44
1:B:301:PRO:HB2	1:B:456:GLY:HA3	1.99	0.43
1:B:306:LYS:H	1:B:306:LYS:CE	2.31	0.43
1:B:75:LEU:HD23	3:B:500:LEH:H13A	2.00	0.43
1:B:190:ARG:HG2	1:B:198:TYR:CZ	2.53	0.43
1:A:75:LEU:CD2	3:A:500:LEH:H13A	2.48	0.43
2:A:470[A]:HEM:HMB2	2:A:470[A]:HEM:HBB2	2.00	0.43
1:A:147:ARG:HG2	1:A:164:SER:HB3	2.01	0.43
1:A:400:CYS:HA	2:A:470[A]:HEM:CHA	2.50	0.42
1:B:106:SER:HB3	1:B:233:LEU:HD23	2.01	0.42
1:B:118[A]:MET:HE2	6:B:579:HOH:O	2.20	0.42
1:A:81:PHE:HB3	1:A:209:ILE:HG12	2.02	0.42
1:B:457:GLY:HA2	6:B:904:HOH:O	2.20	0.42
1:A:189:GLN:O	1:A:190:ARG:HG3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:400:CYS:HA	2:B:471[A]:HEM:C4D	2.56	0.41
1:A:187:LYS:C	1:A:189:GLN:H	2.29	0.40
1:B:141:VAL:HB	1:B:142:PRO:HD3	2.03	0.40
1:B:322:LEU:HD23	6:B:723:HOH:O	2.20	0.40
2:B:471[A]:HEM:C1D	3:B:500:LEH:H19	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/470 (99%)	450 (96%)	15 (3%)	2 (0%)	30	15
1	B	465/470 (99%)	452 (97%)	11 (2%)	2 (0%)	30	15
All	All	932/940 (99%)	902 (97%)	26 (3%)	4 (0%)	30	15

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	230	SER
1	B	266	HIS
1	A	266	HIS
1	B	196	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/411 (100%)	408 (99%)	3 (1%)	76	64
1	B	409/411 (100%)	407 (100%)	2 (0%)	81	72
All	All	820/822 (100%)	815 (99%)	5 (1%)	78	68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	LEU
1	A	230	SER
1	A	455	LEU
1	B	148	LEU
1	B	230	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	70	ASN
1	A	134	ASN
1	A	201	ASN
1	A	359	GLN
1	A	403	GLN
1	B	70	ASN
1	B	73	GLN
1	B	109	GLN
1	B	359	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	471[B]	3,1	50,50,50	1.72	8 (16%)	67,82,82	1.39	12 (17%)
2	HEM	B	471[A]	3,1	50,50,50	1.73	8 (16%)	67,82,82	1.31	6 (8%)
2	HEM	A	470[A]	3,1	50,50,50	1.78	10 (20%)	67,82,82	1.28	11 (16%)
2	HEM	A	470[B]	3,1	50,50,50	1.77	11 (22%)	67,82,82	1.21	6 (8%)
3	LEH	B	500	2	27,27,27	1.83	4 (14%)	32,32,32	1.13	4 (12%)
5	MES	B	490	-	12,12,12	1.86	1 (8%)	15,16,16	5.70	7 (46%)
3	LEH	A	500	2	27,27,27	1.73	4 (14%)	32,32,32	1.04	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	471[B]	3,1	-	2/14/54/54	-
2	HEM	B	471[A]	3,1	-	2/14/54/54	-
2	HEM	A	470[A]	3,1	-	0/14/54/54	-
2	HEM	A	470[B]	3,1	-	3/14/54/54	-
3	LEH	B	500	2	-	13/26/26/26	0/1/1/1
5	MES	B	490	-	-	2/6/14/14	0/1/1/1
3	LEH	A	500	2	-	7/26/26/26	0/1/1/1

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	470[B]	HEM	C3D-C2D	7.55	1.53	1.36
2	B	471[B]	HEM	C3D-C2D	6.91	1.51	1.36
2	A	470[A]	HEM	C3D-C2D	6.90	1.51	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	471[A]	HEM	C3D-C2D	6.57	1.50	1.36
5	B	490	MES	C8-S	-5.91	1.69	1.77
3	B	500	LEH	C20-C21	-5.74	1.22	1.35
3	B	500	LEH	C20-N2	-5.63	1.23	1.37
3	A	500	LEH	C20-C21	-5.37	1.23	1.35
3	A	500	LEH	C20-N2	-5.04	1.25	1.37
2	B	471[B]	HEM	FE-ND	4.78	2.09	1.94
2	B	471[A]	HEM	FE-ND	4.70	2.09	1.94
2	A	470[A]	HEM	FE-ND	4.01	2.07	1.94
2	B	471[A]	HEM	FE-NC	3.63	2.07	1.95
2	A	470[A]	HEM	FE-NA	3.55	2.06	1.95
2	A	470[A]	HEM	CAC-C3C	3.45	1.56	1.47
2	A	470[A]	HEM	FE-NB	3.38	2.05	1.94
3	A	500	LEH	C21-N3	-3.31	1.25	1.37
2	A	470[B]	HEM	FE-ND	3.26	2.04	1.94
2	B	471[B]	HEM	FE-NC	3.17	2.05	1.95
2	B	471[A]	HEM	CAC-C3C	3.10	1.55	1.47
2	B	471[B]	HEM	CMD-C2D	3.09	1.57	1.50
2	B	471[A]	HEM	CMA-C3A	3.08	1.57	1.50
3	B	500	LEH	C21-N3	-3.04	1.26	1.37
2	B	471[A]	HEM	CAB-C3B	3.03	1.55	1.47
2	A	470[B]	HEM	CAB-C3B	3.03	1.55	1.47
2	A	470[B]	HEM	FE-NA	3.02	2.05	1.95
2	A	470[B]	HEM	FE-NB	2.98	2.04	1.94
3	A	500	LEH	C19-N2	-2.90	1.27	1.35
2	B	471[B]	HEM	CAB-C3B	2.83	1.54	1.47
2	A	470[A]	HEM	CAB-C3B	2.80	1.54	1.47
2	B	471[B]	HEM	CAC-C3C	2.75	1.54	1.47
2	A	470[B]	HEM	CAC-C3C	2.68	1.54	1.47
3	B	500	LEH	C19-N2	-2.66	1.28	1.35
2	A	470[B]	HEM	C2A-C3A	-2.51	1.32	1.38
2	B	471[A]	HEM	C2A-C3A	-2.47	1.32	1.38
2	A	470[B]	HEM	CMB-C2B	2.47	1.55	1.50
2	B	471[B]	HEM	C2A-C3A	-2.38	1.32	1.38
2	A	470[A]	HEM	FE-NC	2.32	2.02	1.95
2	B	471[A]	HEM	FE-NA	2.31	2.02	1.95
2	A	470[A]	HEM	O1A-CGA	2.26	1.29	1.22
2	A	470[B]	HEM	O1D-CGD	2.19	1.29	1.22
2	B	471[B]	HEM	CMB-C2B	2.18	1.55	1.50
2	A	470[A]	HEM	C2A-C3A	-2.12	1.33	1.38
2	A	470[A]	HEM	CMC-C2C	2.05	1.55	1.50
2	A	470[B]	HEM	CMD-C2D	2.01	1.54	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	470[B]	HEM	FE-NC	2.01	2.01	1.95

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	490	MES	O1S-S-C8	-12.50	87.84	106.73
5	B	490	MES	O2S-S-C8	-11.49	89.37	106.73
5	B	490	MES	O3S-S-C8	-10.26	85.94	106.00
5	B	490	MES	C5-N4-C3	5.80	121.34	108.84
5	B	490	MES	O3S-S-O1S	5.79	125.87	111.40
2	A	470[B]	HEM	C4D-ND-C1D	3.95	109.88	105.21
5	B	490	MES	C7-N4-C5	3.93	121.71	111.24
2	B	471[B]	HEM	C4D-ND-C1D	3.86	109.78	105.21
2	A	470[A]	HEM	C4D-ND-C1D	3.84	109.76	105.21
2	B	471[A]	HEM	CHD-C1D-ND	3.52	128.22	124.42
3	B	500	LEH	C6-N1-C7	3.38	130.16	121.68
2	B	471[B]	HEM	CHC-C4B-NB	3.30	127.97	124.42
2	B	471[A]	HEM	CAD-C3D-C4D	3.14	130.17	124.70
2	B	471[A]	HEM	C4D-ND-C1D	3.12	108.90	105.21
2	A	470[B]	HEM	CAD-C3D-C4D	3.07	130.04	124.70
2	B	471[B]	HEM	C3B-C4B-NB	-2.91	107.38	109.47
2	B	471[A]	HEM	C3B-C2B-C1B	2.82	108.53	106.41
2	B	471[B]	HEM	C3B-C2B-C1B	2.66	108.41	106.41
2	B	471[B]	HEM	C1B-NB-C4B	2.66	108.36	105.21
2	A	470[A]	HEM	CHD-C1D-ND	2.65	127.27	124.42
2	A	470[A]	HEM	CAD-C3D-C4D	2.50	129.05	124.70
2	B	471[A]	HEM	CBD-CAD-C3D	-2.49	105.65	112.53
3	A	500	LEH	C5-C6-N1	-2.48	104.99	110.58
2	B	471[A]	HEM	C2A-C1A-NA	-2.46	107.42	110.15
2	B	471[B]	HEM	CHD-C4C-NC	2.46	127.13	124.45
2	B	471[B]	HEM	CMD-C2D-C1D	2.42	128.82	125.03
2	A	470[A]	HEM	C1B-NB-C4B	2.39	108.03	105.21
2	A	470[B]	HEM	CHD-C4C-NC	2.37	127.03	124.45
2	B	471[B]	HEM	CAD-C3D-C4D	2.35	128.79	124.70
2	A	470[A]	HEM	C3B-C2B-C1B	2.34	108.17	106.41
2	A	470[A]	HEM	CBA-CAA-C2A	-2.31	106.13	112.53
2	A	470[A]	HEM	CHC-C4B-NB	2.29	126.89	124.42
2	B	471[B]	HEM	CHB-C1B-NB	2.28	127.19	124.37
3	A	500	LEH	N2-C19-N3	-2.26	106.07	111.45
2	A	470[B]	HEM	CBA-CAA-C2A	-2.20	106.44	112.53
2	B	471[B]	HEM	C1D-C2D-C3D	-2.20	104.67	106.98
2	A	470[B]	HEM	CHC-C4B-NB	2.19	126.78	124.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	470[A]	HEM	O1A-CGA-CBA	-2.16	116.25	123.09
2	B	471[B]	HEM	C2B-C1B-NB	-2.15	107.37	109.84
2	A	470[A]	HEM	O2A-CGA-CBA	2.11	120.67	114.00
2	B	471[B]	HEM	CHA-C4D-ND	2.10	126.97	124.37
3	B	500	LEH	C5-C6-N1	-2.09	105.87	110.58
5	B	490	MES	C7-N4-C3	2.08	116.79	111.24
3	B	500	LEH	N2-C19-N3	-2.07	106.51	111.45
2	A	470[B]	HEM	CHD-C1D-ND	2.02	126.60	124.42
3	B	500	LEH	O2-C1-O1	-2.02	119.50	124.08
2	A	470[A]	HEM	C2B-C1B-NB	-2.02	107.52	109.84
2	A	470[A]	HEM	CBD-CAD-C3D	-2.01	106.97	112.53

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500	LEH	C7-C8-C9-C10
3	B	500	LEH	C7-C8-C9-C10
3	B	500	LEH	C3-C4-C5-C6
3	B	500	LEH	C9-C10-C11-C12
3	B	500	LEH	C12-C13-C14-C15
3	B	500	LEH	C10-C11-C12-C13
3	A	500	LEH	C11-C10-C9-C8
3	A	500	LEH	C9-C10-C11-C12
3	B	500	LEH	C2-C4-C5-C6
3	A	500	LEH	C12-C13-C14-C15
5	B	490	MES	C8-C7-N4-C3
3	B	500	LEH	C11-C10-C9-C8
3	B	500	LEH	O3-C7-C8-C9
3	B	500	LEH	N1-C7-C8-C9
5	B	490	MES	C8-C7-N4-C5
3	B	500	LEH	C14-C15-C16-C17
3	B	500	LEH	C17-C18-N2-C20
2	B	471[B]	HEM	CAA-CBA-CGA-O1A
2	B	471[B]	HEM	CAA-CBA-CGA-O2A
3	B	500	LEH	C17-C18-N2-C19
2	A	470[B]	HEM	CAA-CBA-CGA-O2A
3	B	500	LEH	C13-C14-C15-C16
2	A	470[B]	HEM	CAA-CBA-CGA-O1A
3	A	500	LEH	O3-C7-C8-C9
2	B	471[A]	HEM	CAD-CBD-CGD-O1D
3	A	500	LEH	N1-C7-C8-C9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	500	LEH	C14-C15-C16-C17
2	B	471[A]	HEM	CAD-CBD-CGD-O2D
2	A	470[B]	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.

5 monomers are involved in 20 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	471[B]	HEM	1	0
2	B	471[A]	HEM	5	0
2	A	470[A]	HEM	3	0
3	B	500	LEH	5	0
3	A	500	LEH	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEM B 471 (B)




Ligand HEM B 471 (A)




Ligand HEM A 470 (A)




Ligand HEM A 470 (B)



Ligand LEH B 500





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/470 (96%)	-0.02	17 (3%) 45 49	5, 15, 31, 40	15 (3%)
1	B	457/470 (97%)	0.03	36 (7%) 18 21	5, 14, 35, 46	10 (2%)
All	All	911/940 (96%)	0.01	53 (5%) 29 32	5, 14, 33, 46	25 (2%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	220	ILE	4.5
1	B	225	ALA	4.4
1	B	221	ALA	3.9
1	B	216	VAL	3.8
1	A	457	GLY	3.7
1	B	191	ALA	3.7
1	B	219	ILE	3.4
1	A	4	GLU	3.4
1	A	196	PRO	3.4
1	B	224	LYS	3.2
1	B	215	LEU	3.2
1	A	189	GLN	3.2
1	B	197	ALA	3.1
1	B	174	ILE	3.1
1	A	230	SER	3.0
1	A	220	ILE	2.9
1	B	171	HIS	2.9
1	A	191	ALA	2.9
1	A	198	TYR	2.9
1	B	198	TYR	2.9
1	B	2	ILE	2.9
1	B	457	GLY	2.8
1	B	196	PRO	2.7
1	A	188	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	GLN	2.6
1	B	1	THR	2.6
1	B	169	GLN	2.6
1	B	188	LEU	2.6
1	B	218	LYS	2.6
1	A	221	ALA	2.5
1	A	231	ASP	2.5
1	B	172	PRO	2.5
1	A	194	ASP	2.4
1	B	223	ARG	2.4
1	B	211	VAL	2.4
1	B	227	GLY	2.3
1	B	264	ALA	2.3
1	B	228	GLU	2.3
1	A	219	ILE	2.3
1	B	173	PHE	2.2
1	B	184	ALA	2.2
1	B	175	THR	2.2
1	A	228	GLU	2.1
1	B	193	PRO	2.1
1	B	455	LEU	2.1
1	A	369	ASP	2.1
1	B	222	ASP	2.1
1	B	192	ASN	2.0
1	B	203	ARG	2.0
1	B	268	THR	2.0
1	A	229	GLN	2.0
1	B	199	ASP	2.0
1	A	216	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	LEH	B	500	27/27	0.74	0.19	17,50,58,59	0
3	LEH	A	500	27/27	0.80	0.19	15,50,55,56	0
5	MES	B	490	12/12	0.82	0.18	57,58,60,61	0
4	MG	B	470	1/1	0.94	0.17	38,38,38,38	0
2	HEM	A	470[A]	43/43	0.98	0.05	10,11,13,16	43
2	HEM	A	470[B]	43/43	0.98	0.05	8,9,11,12	43
2	HEM	B	471[A]	43/43	0.98	0.05	7,11,12,16	43
2	HEM	B	471[B]	43/43	0.98	0.05	10,12,14,16	43

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around LEH A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 470 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around HEM A 470 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM B 471 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 471 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.