



wwPDB EM Validation Summary Report ⓘ

Aug 6, 2026 – 04:48 AM JST

PDB ID : 7D58 / pdb_00007d58
EMDB ID : EMD-30577
Title : cryo-EM structure of human RNA polymerase III in elongating state
Authors : Wang, Q.; Wan, F.; Lan, P.; Wu, J.; Lei, M.
Deposited on : 2020-09-25
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	13054 (2.40 - 3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1390	
2	B	1133	
3	C	346	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	148	
5	E	210	
6	F	127	
7	G	204	
8	H	150	
9	I	108	
10	J	67	
11	K	133	
12	L	58	
13	M	708	
14	N	398	
15	O	534	
16	P	316	
17	Q	223	
18	R	41	
19	S	41	
20	T	20	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
22	SF4	P	400	-	-	X	-

2 Entry composition

There are 22 unique types of molecules in this entry. The entry contains 40568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase III subunit RPC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1387	Total	C	N	O	S	0	0
			10886	6899	1898	2016	73		

- Molecule 2 is a protein called DNA-directed RNA polymerase III subunit RPC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1095	Total	C	N	O	S	0	0
			8668	5492	1513	1595	68		

- Molecule 3 is a protein called DNA-directed RNA polymerases I and III subunit RPAC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	345	Total	C	N	O	S	0	0
			2752	1732	490	518	12		

- Molecule 4 is a protein called DNA-directed RNA polymerase III subunit RPC9.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	126	Total	C	N	O	S	0	0
			1013	634	176	200	3		

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	210	Total	C	N	O	S	0	0
			1723	1088	301	325	9		

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	79	Total	C	N	O	S	0	0
			636	407	108	116	5		

- Molecule 7 is a protein called DNA-directed RNA polymerase III subunit RPC8.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	187	Total	C	N	O	S	0	0
			1493	964	235	287	7		

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	149	Total	C	N	O	S	0	0
			1197	759	195	238	5		

- Molecule 9 is a protein called DNA-directed RNA polymerase III subunit RPC10.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	108	Total	C	N	O	S	0	0
			857	529	158	157	13		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	67	Total	C	N	O	S	0	0
			533	345	90	92	6		

- Molecule 11 is a protein called DNA-directed RNA polymerases I and III subunit RPAC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	107	Total	C	N	O	S	0	0
			856	531	153	165	7		

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	47	Total	C	N	O	S	0	0
			397	246	77	68	6		

- Molecule 13 is a protein called DNA-directed RNA polymerase III subunit RPC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	222	Total	C	N	O	S	0	0
			1797	1134	307	346	10		

- Molecule 14 is a protein called DNA-directed RNA polymerase III subunit RPC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	136	Total	C	N	O	S	0	0
			1038	653	188	192	5		

- Molecule 15 is a protein called DNA-directed RNA polymerase III subunit RPC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	490	Total	C	N	O	S	0	0
			3913	2468	678	742	25		

- Molecule 16 is a protein called DNA-directed RNA polymerase III subunit RPC6.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	139	Total	C	N	O	S	0	0
			1103	700	178	214	11		

- Molecule 17 is a protein called DNA-directed RNA polymerase III subunit RPC7.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	86	Total	C	N	O	S	0	0
			746	482	125	133	6		

- Molecule 18 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	17	Total	C	N	O	P	0	0
			346	165	63	101	17		

- Molecule 19 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	S	24	Total	C	N	O	P	0	0
			493	234	90	145	24		


- Molecule 20 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	T	5	Total	C	N	O	P	0	0
			106	47	17	37	5		

- Molecule 21 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
21	A	2	Total 2	Zn 2	0
21	B	1	Total 1	Zn 1	0
21	I	2	Total 2	Zn 2	0
21	J	1	Total 1	Zn 1	0
21	L	1	Total 1	Zn 1	0

- Molecule 22 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			AltConf
22	P	1	Total 8	Fe 4	S 4	0

3 Residue-property plots



These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase III subunit RPC1







• Molecule 2: DNA-directed RNA polymerase III subunit RPC2







- Molecule 3: DNA-directed RNA polymerases I and III subunit RPAC1




- Molecule 4: DNA-directed RNA polymerase III subunit RPC9




- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2




- Molecule 7: DNA-directed RNA polymerase III subunit RPC8




- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3





- Molecule 9: DNA-directed RNA polymerase III subunit RPC10






- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5








- Molecule 11: DNA-directed RNA polymerases I and III subunit RPAC2





- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 13: DNA-directed RNA polymerase III subunit RPC5








• Molecule 16: DNA-directed RNA polymerase III subunit RPC6



• Molecule 17: DNA-directed RNA polymerase III subunit RPC7



DA	DT	DT	DT	DA	DG	DC	DC	DA	DA	DT	DG	DG	DG	DG	DA	TS4	G55	A56	C57	AS8	C59	G60	C61	AS2	AG3	C70
----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

[illegible]

C	A	U	A	U	G	A	C	U	U	G	U	A	A	C	G4	U5	G6	U7	C8
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	----	----	----	----	----

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	155030	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.082	Depositor
Minimum map value	-0.031	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.004	Depositor
Map size ($\text{\AA}$)	352.0, 352.0, 352.0	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1, 1.1, 1.1	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/11083	0.43	3/14946 (0.0%)
2	B	0.14	0/8833	0.40	0/11912
3	C	0.15	0/2806	0.41	0/3803
4	D	0.14	0/1026	0.43	0/1384
5	E	0.15	0/1753	0.46	0/2368
6	F	0.13	0/646	0.37	0/873
7	G	0.15	0/1534	0.42	0/2087
8	H	0.13	0/1219	0.38	0/1644
9	I	0.17	0/878	0.48	0/1186
10	J	0.12	0/542	0.35	0/730
11	K	0.12	0/871	0.36	0/1174
12	L	0.14	0/403	0.40	0/536
13	M	0.15	0/1837	0.38	0/2481
14	N	0.17	0/1051	0.43	0/1417
15	O	0.14	0/3972	0.41	0/5359
16	P	0.17	0/1126	0.45	0/1522
17	Q	0.16	0/769	0.45	0/1038
18	R	0.13	0/387	0.33	0/594
19	S	0.14	0/552	0.29	0/850
20	T	0.04	0/117	0.13	0/180
All	All	0.14	0/41405	0.41	3/56084 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
4	D	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
15	O	0	1
All	All	0	4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1180	ASN	N-CA-C	5.07	116.23	108.52
1	A	914	GLY	CA-C-N	5.04	131.16	121.54
1	A	914	GLY	C-N-CA	5.04	131.16	121.54

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	589	LEU	Peptide
2	B	751	ILE	Peptide
4	D	77	LYS	Peptide
15	O	68	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10886	0	11135	194	0
2	B	8668	0	8796	147	0
3	C	2752	0	2725	50	0
4	D	1013	0	1040	22	0
5	E	1723	0	1745	24	0
6	F	636	0	667	10	0
7	G	1493	0	1429	33	0
8	H	1197	0	1156	22	0
9	I	857	0	815	26	0
10	J	533	0	555	5	0
11	K	856	0	840	23	0
12	L	397	0	404	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	M	1797	0	1756	42	0
14	N	1038	0	1092	30	0
15	O	3913	0	3988	76	0
16	P	1103	0	1067	44	0
17	Q	746	0	749	15	0
18	R	346	0	192	4	0
19	S	493	0	271	12	0
20	T	106	0	54	3	0
21	A	2	0	0	0	0
21	B	1	0	0	0	0
21	I	2	0	0	0	0
21	J	1	0	0	0	0
21	L	1	0	0	0	0
22	P	8	0	0	3	0
All	All	40568	0	40476	678	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 678 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:G:92:CYS:HB2	7:G:125:GLN:HB3	1.55	0.89
7:G:96:GLY:HA3	7:G:109:ILE:O	1.73	0.88
9:I:35:THR:HG21	14:N:29:ARG:HH22	1.41	0.85
1:A:995:THR:HG22	1:A:997:PRO:HD2	1.61	0.81
7:G:99:VAL:HG13	7:G:107:ILE:HG23	1.62	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1385/1390 (100%)	1304 (94%)	80 (6%)	1 (0%)	48	77
2	B	1089/1133 (96%)	1030 (95%)	59 (5%)	0	100	100
3	C	343/346 (99%)	316 (92%)	27 (8%)	0	100	100
4	D	124/148 (84%)	114 (92%)	9 (7%)	1 (1%)	16	44
5	E	208/210 (99%)	192 (92%)	16 (8%)	0	100	100
6	F	77/127 (61%)	73 (95%)	4 (5%)	0	100	100
7	G	181/204 (89%)	167 (92%)	14 (8%)	0	100	100
8	H	147/150 (98%)	138 (94%)	9 (6%)	0	100	100
9	I	106/108 (98%)	98 (92%)	8 (8%)	0	100	100
10	J	65/67 (97%)	63 (97%)	2 (3%)	0	100	100
11	K	105/133 (79%)	104 (99%)	1 (1%)	0	100	100
12	L	45/58 (78%)	42 (93%)	3 (7%)	0	100	100
13	M	214/708 (30%)	206 (96%)	8 (4%)	0	100	100
14	N	130/398 (33%)	117 (90%)	13 (10%)	0	100	100
15	O	484/534 (91%)	468 (97%)	16 (3%)	0	100	100
16	P	135/316 (43%)	127 (94%)	8 (6%)	0	100	100
17	Q	84/223 (38%)	77 (92%)	7 (8%)	0	100	100
All	All	4922/6253 (79%)	4636 (94%)	284 (6%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	78	LEU
1	A	591	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1208/1212 (100%)	1207 (100%)	1 (0%)	88	97

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	958/988 (97%)	957 (100%)	1 (0%)	88	97
3	C	301/302 (100%)	301 (100%)	0	100	100
4	D	117/136 (86%)	117 (100%)	0	100	100
5	E	192/192 (100%)	192 (100%)	0	100	100
6	F	69/111 (62%)	69 (100%)	0	100	100
7	G	167/181 (92%)	166 (99%)	1 (1%)	78	93
8	H	130/131 (99%)	130 (100%)	0	100	100
9	I	94/94 (100%)	94 (100%)	0	100	100
10	J	56/56 (100%)	56 (100%)	0	100	100
11	K	96/119 (81%)	96 (100%)	0	100	100
12	L	44/55 (80%)	44 (100%)	0	100	100
13	M	200/622 (32%)	200 (100%)	0	100	100
14	N	119/347 (34%)	119 (100%)	0	100	100
15	O	439/476 (92%)	438 (100%)	1 (0%)	87	96
16	P	125/280 (45%)	125 (100%)	0	100	100
17	Q	83/195 (43%)	83 (100%)	0	100	100
All	All	4398/5497 (80%)	4394 (100%)	4 (0%)	87	97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1153	ILE
2	B	333	ILE
7	G	97	VAL
15	O	462	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 97 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	172	HIS
9	I	10	ASN
3	C	265	GLN
4	D	88	ASN
11	K	45	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
20	T	4/20 (20%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
22	SF4	P	400	16	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
22	SF4	P	400	16	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
22	P	400	SF4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-30577. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 183



Y Index: 164




Z Index: 141


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.004. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 658 nm^3 ; this corresponds to an approximate mass of 594 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-30577 and PDB model 7D58. Per-residue inclusion information can be found in [section 3](#) on [page 8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.004 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.004).

9.4 Atom inclusion ⓘ



At the recommended contour level, 79% of all backbone atoms, 67% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.004) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6710	 0.0140
A	 0.6110	 -0.0100
B	 0.6330	 0.0030
C	 0.7690	 0.0600
D	 0.7950	 0.0590
E	 0.6680	 -0.0180
F	 0.6430	 -0.0020
G	 0.7560	 0.0490
H	 0.6730	 -0.0020
I	 0.6180	 0.0040
J	 0.6910	 0.0220
K	 0.7340	 0.0670
L	 0.7770	 0.0640
M	 0.7330	 0.0330
N	 0.7340	 0.0040
O	 0.7060	 0.0240
P	 0.7590	 0.0380
Q	 0.6400	 0.0300
R	 0.8560	 0.1390
S	 0.6900	 0.0710
T	 0.4430	 0.0250

