



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 10:06 PM JST

PDB ID : 7D9F / pdb_00007d9f
Title : SpdH Spermidine dehydrogenase SeMet Structure
Authors : Che, S.; Zhang, Q.; Bartlam, M.
Deposited on : 2020-10-13
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 10713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spermidine dehydrogenase, SpdH.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	591	Total	C	N	O	S	Se	0	0	0
			4649	2942	829	861	2	15			
1	B	587	Total	C	N	O	S	Se	0	0	0
			4629	2930	825	857	2	15			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



- GOL
-
- The diagram shows the skeletal structure of 1,2,3-propanetriol (glycerol). It consists of a three-carbon chain. The first carbon (C1) is bonded to a hydroxyl group (HO, O1). The second carbon (C2) is bonded to two hydroxyl groups (OH, O2). The third carbon (C3) is bonded to a hydroxyl group (OH, O3). The labels C1, C2, C3, O1, O2, and O3 are in green, while the labels HO, OH, and OH are in red.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	578	Total	O	0	0
			578	578		
6	B	630	Total	O	0	0
			630	630		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Spermidine dehydrogenase, SpdH



- Molecule 1: Spermidine dehydrogenase, SpdH



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.20Å 85.59Å 99.62Å 90.00° 98.33° 90.00°	Depositor
Resolution (Å)	39.25 – 1.85 39.25 – 1.85	Depositor EDS
% Data completeness (in resolution range)	84.0 (39.25-1.85) 90.4 (39.25-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.85Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.150 , 0.190 0.150 , 0.191	Depositor DCC
R_{free} test set	2000 reflections (2.16%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10713	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD, SO4, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/4752	0.48	0/6417
1	B	0.30	0/4731	0.49	0/6388
All	All	0.29	0/9483	0.49	0/12805

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4649	0	4520	28	1
1	B	4629	0	4502	26	0
2	A	53	0	31	1	0
2	B	53	0	31	1	0
3	A	43	0	30	6	0
3	B	43	0	30	5	0
4	A	12	0	16	0	0
4	B	18	0	24	1	0
5	A	5	0	0	0	0
6	A	578	0	0	7	1
6	B	630	0	0	7	2
All	All	10713	0	9184	57	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319:GLU:OE1	6:A:801:HOH:O	2.13	0.67
1:B:238:ARG:NH1	6:B:803:HOH:O	2.27	0.67
1:A:449:ARG:NH2	6:A:805:HOH:O	2.29	0.65
1:B:107:HIS:CE1	3:B:703:HEM:HBB1	2.33	0.63
1:B:449:ARG:NH1	6:B:807:HOH:O	2.33	0.61

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:GLU:OE2	1:A:395:TYR:OH[2_545]	2.04	0.16
6:A:1135:HOH:O	6:B:1376:HOH:O[2_646]	2.14	0.06
6:B:1353:HOH:O	6:B:1361:HOH:O[2_546]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/620 (95%)	575 (98%)	14 (2%)	0	100	100
1	B	585/620 (94%)	574 (98%)	11 (2%)	0	100	100
All	All	1174/1240 (95%)	1149 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/480 (99%)	468 (98%)	7 (2%)	57	47
1	B	474/480 (99%)	471 (99%)	3 (1%)	78	73
All	All	949/960 (99%)	939 (99%)	10 (1%)	65	57

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	198	MSE
1	B	416	MSE
1	B	575	GLU
1	A	270	GLU
1	A	286	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	452	GLN
1	B	281	ASN
1	B	505	ASN
1	B	317	GLN
1	A	281	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	702	1	50,50,50	1.40	8 (16%)	66,82,82	1.21	8 (12%)
4	GOL	B	704	-	5,5,5	1.05	0	5,5,5	0.73	0
4	GOL	A	705	-	5,5,5	0.78	0	5,5,5	1.14	1 (20%)
5	SO4	A	704	-	4,4,4	0.14	0	6,6,6	0.13	0
4	GOL	A	703	-	5,5,5	1.15	0	5,5,5	0.68	0
2	FAD	B	702	-	56,58,58	0.47	1 (1%)	81,89,89	0.55	1 (1%)
4	GOL	B	705	-	5,5,5	0.72	0	5,5,5	1.12	1 (20%)
4	GOL	B	701	-	5,5,5	0.98	0	5,5,5	0.87	0
2	FAD	A	701	-	56,58,58	0.44	0	81,89,89	0.52	0
3	HEM	B	703	1	50,50,50	1.44	9 (18%)	66,82,82	1.36	11 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	702	1	-	2/14/54/54	-
4	GOL	B	704	-	-	2/4/4/4	-
4	GOL	A	705	-	-	2/4/4/4	-
4	GOL	A	703	-	-	3/4/4/4	-
2	FAD	B	702	-	-	5/34/50/50	0/6/6/6
4	GOL	B	705	-	-	2/4/4/4	-
4	GOL	B	701	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	701	-	-	5/34/50/50	0/6/6/6
3	HEM	B	703	1	-	2/14/54/54	-

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	HEM	FE-NC	4.15	2.09	1.95
3	B	703	HEM	FE-NC	4.03	2.08	1.95
3	B	703	HEM	FE-NB	3.23	2.04	1.94
3	B	703	HEM	FE-NA	3.09	2.05	1.95
3	B	703	HEM	CAB-C3B	2.96	1.55	1.47

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	703	HEM	C4A-NA-C1A	3.64	108.91	105.35
3	A	702	HEM	C4D-ND-C1D	3.26	108.44	105.07
3	B	703	HEM	C1B-NB-C4B	3.17	108.35	105.07
3	B	703	HEM	C4D-ND-C1D	3.13	108.31	105.07
3	B	703	HEM	C4C-NC-C1C	3.01	108.29	105.35

There are no chirality outliers.

5 of 25 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	FAD	N10-C1'-C2'-O2'
2	A	701	FAD	N10-C1'-C2'-C3'
2	B	702	FAD	N10-C1'-C2'-O2'
2	B	702	FAD	N10-C1'-C2'-C3'
4	A	705	GOL	O1-C1-C2-C3


There are no ring outliers.


5 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	HEM	6	0
2	B	702	FAD	1	0
4	B	701	GOL	1	0
2	A	701	FAD	1	0
3	B	703	HEM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/620 (92%)	-0.41	7 (1%) 76 80	11, 20, 35, 65	0
1	B	572/620 (92%)	-0.50	0 100 100	11, 18, 32, 44	0
All	All	1148/1240 (92%)	-0.45	7 (0%) 85 88	11, 19, 34, 65	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	29	ALA	2.9
1	A	314	GLU	2.8
1	A	313	SER	2.7
1	A	30	PRO	2.5
1	A	312	PRO	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	A	704	5/5	0.63	0.13	51,52,59,72	0
4	GOL	B	705	6/6	0.78	0.14	26,30,32,38	0
4	GOL	A	705	6/6	0.79	0.13	28,32,37,41	0
4	GOL	A	703	6/6	0.89	0.10	27,31,31,32	0
4	GOL	B	704	6/6	0.93	0.09	23,24,29,30	0
4	GOL	B	701	6/6	0.94	0.07	21,26,29,30	0
3	HEM	A	702	43/43	0.98	0.06	15,19,24,26	0
3	HEM	B	703	43/43	0.98	0.06	14,19,26,29	0
2	FAD	A	701	53/53	0.98	0.04	8,13,16,17	0
2	FAD	B	702	53/53	0.98	0.04	10,11,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM B 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.