wwPDB X-ray Structure Validation Summary Report (i)

Aug 5, 2026 – 03:11 PM EDT

PDB ID	:	9DQ2 / pdb_00009dq2
Title	:	Crystal structure of HrmJ from Streptomyces sp. CFMR 7 (HrmJ-ssc) complexed with vanadyl(IV)-oxo, succinate and 6-nitronorleucine
Authors	:	Zheng, Y.-C.; Chang, W.-C.
Deposited on	:	2024-09-23
Resolution	:	1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	6% 88% 8% .
1	B	236	8% 89% 7% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	BR	A	308	-	-	X	-
6	BR	B	307	-	-	X	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 4410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BsmA domain containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	7	0
			1826	1157	321	344	4			
1	B	226	Total	C	N	O	S	0	9	0
			1847	1172	326	345	4			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	231	HIS	-	expression tag	UNP A0A0M5J3M0
A	232	HIS	-	expression tag	UNP A0A0M5J3M0
A	233	HIS	-	expression tag	UNP A0A0M5J3M0
A	234	HIS	-	expression tag	UNP A0A0M5J3M0
A	235	HIS	-	expression tag	UNP A0A0M5J3M0
A	236	HIS	-	expression tag	UNP A0A0M5J3M0
B	231	HIS	-	expression tag	UNP A0A0M5J3M0
B	232	HIS	-	expression tag	UNP A0A0M5J3M0
B	233	HIS	-	expression tag	UNP A0A0M5J3M0
B	234	HIS	-	expression tag	UNP A0A0M5J3M0
B	235	HIS	-	expression tag	UNP A0A0M5J3M0
B	236	HIS	-	expression tag	UNP A0A0M5J3M0

- Molecule 2 is 6-nitro-L-norleucine (CCD ID: 6HN) (formula: $C_6H_{12}N_2O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 12 6 2 4	0	0
2	B	1	Total C N O 12 6 2 4	0	0

- Molecule 3 is oxovanadium(2+) (CCD ID: VVO) (formula: OV) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O V 2 1 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O V 2 1 1	0	0

- Molecule 4 is SUCCINIC ACID (CCD ID: SIN) (formula: $C_4H_6O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 4 4	0	0
4	B	1	Total C O 8 4 4	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total Br 3 3	0	0
6	B	1	Total Br 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	337	Total O 337 337	0	0
7	B	322	Total O 322 322	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BsmA domain containing protein

Chain A:






• Molecule 1: BsmA domain containing protein

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	75.11Å 78.84Å 97.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.36 – 1.55 41.36 – 1.55	Depositor EDS
% Data completeness (in resolution range)	96.9 (41.36-1.55) 95.3 (41.36-1.55)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$< I/\sigma(I) >$ [1]	2.22 (at 1.55Å)	Xtriage
Refinement program	PHENIX 1.21	Depositor
R, R_{free}	0.196 , 0.211 0.195 , 0.210	Depositor DCC
R_{free} test set	4007 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å^2)	12.1	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 39.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4410	wwPDB-VP
Average B, all atoms (Å^2)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.6645e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIN, GOL, 6HN, VVO, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.44	0/1887	0.66	1/2569 (0.0%)
1	B	0.47	0/1908	0.65	0/2597
All	All	0.45	0/3795	0.65	1/5166 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	145	LEU	CA-CB-CG	-5.67	96.44	116.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	201	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1826	0	1755	17	0
1	B	1847	0	1784	17	0
2	A	12	0	11	0	0
2	B	12	0	11	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	8	0	4	0	0
4	B	8	0	4	0	0
5	A	12	0	16	0	0
5	B	18	0	24	0	0
6	A	3	0	0	3	1
6	B	1	0	0	3	0
7	A	337	0	0	14	3
7	B	322	0	0	13	5
All	All	4410	0	3609	37	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:307:BR:BR	7:B:651:HOH:O	2.04	1.30
1:A:79[B]:ARG:NH2	7:A:401:HOH:O	1.65	1.25
1:A:6:GLU:OE1	7:A:402:HOH:O	1.66	1.10
1:B:6:GLU:OE1	7:B:401:HOH:O	1.74	1.04
1:A:3:LEU:O	7:A:403:HOH:O	1.78	1.00

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:415:HOH:O	7:B:508:HOH:O[3_554]	1.85	0.35
7:A:660:HOH:O	7:B:691:HOH:O[2_555]	1.92	0.28
7:A:586:HOH:O	7:B:537:HOH:O[2_455]	2.10	0.10
7:A:696:HOH:O	7:B:648:HOH:O[2_555]	2.13	0.07
6:A:308:BR:BR	7:B:636:HOH:O[2_455]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/236 (98%)	227 (98%)	4 (2%)	0	100	100
1	B	233/236 (99%)	229 (98%)	4 (2%)	0	100	100
All	All	464/472 (98%)	456 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/195 (99%)	192 (100%)	1 (0%)	81	68
1	B	195/195 (100%)	195 (100%)	0	100	100
All	All	388/390 (100%)	387 (100%)	1 (0%)	86	77

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	145	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	44	GLN
1	B	115	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	6HN	B	301	-	8,11,11	0.86	0	6,13,13	0.88	0
4	SIN	A	303	3	7,7,7	1.00	0	8,8,8	1.84	2 (25%)
5	GOL	A	304	-	5,5,5	0.30	0	5,5,5	0.47	0
5	GOL	A	305	-	5,5,5	0.32	0	5,5,5	0.57	0
3	VVO	B	302	4,1,7	0,1,1	-	-	-		
2	6HN	A	301	-	8,11,11	0.88	0	6,13,13	0.93	0
4	SIN	B	303	3	7,7,7	0.96	0	8,8,8	1.88	2 (25%)
5	GOL	B	305	-	5,5,5	0.20	0	5,5,5	0.95	0
5	GOL	B	306	-	5,5,5	0.38	0	5,5,5	0.28	0
3	VVO	A	302	4,1,7	0,1,1	-	-	-		
5	GOL	B	304	-	5,5,5	0.29	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	304	-	-	0/4/4/4	-
2	6HN	B	301	-	-	1/10/11/11	-
4	SIN	A	303	3	-	4/5/5/5	-
5	GOL	A	305	-	-	4/4/4/4	-
4	SIN	B	303	3	-	2/5/5/5	-
2	6HN	A	301	-	-	1/10/11/11	-
5	GOL	B	305	-	-	4/4/4/4	-
5	GOL	B	306	-	-	0/4/4/4	-
5	GOL	B	304	-	-	0/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	B	303	SIN	C2-C3-C4	-3.39	104.67	113.67
4	A	303	SIN	C3-C2-C1	-3.04	105.61	113.67
4	A	303	SIN	O2-C1-C2	2.47	121.80	114.00
4	B	303	SIN	O4-C4-C3	2.15	120.80	114.00

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	6HN	CG-CD-CE-N1
2	B	301	6HN	CG-CD-CE-N1
5	A	305	GOL	C1-C2-C3-O3
5	B	305	GOL	C1-C2-C3-O3
5	B	305	GOL	O2-C2-C3-O3


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 6HN B 301




Bond lengths



Bond angles




Torsions




Rings


Ligand SIN A 303




Bond lengths



Bond angles




Torsions




Rings


Ligand VVO B 302




Bond lengths



Bond angles




Torsions




Rings


Ligand 6HN A 301




Bond lengths



Bond angles




Torsions




Rings

Ligand SIN B 303



Ligand VVO A 302



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	226/236 (95%)	0.42	15 (6%)	24	29	5, 14, 34, 68	7 (3%)
1	B	226/236 (95%)	0.45	18 (7%)	18	21	5, 13, 32, 70	9 (3%)
All	All	452/472 (95%)	0.43	33 (7%)	21	24	5, 13, 34, 70	16 (3%)

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	50	ALA	6.8
1	A	50	ALA	6.1
1	B	47[A]	LEU	5.2
1	A	52	ASP	4.9
1	A	51	GLU	4.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GOL	B	304	6/6	0.68	0.28	39,51,57,62	0
5	GOL	A	305	6/6	0.77	0.21	42,43,47,51	0
5	GOL	A	304	6/6	0.77	0.17	29,44,49,50	0
5	GOL	B	306	6/6	0.80	0.17	34,38,42,47	0
5	GOL	B	305	6/6	0.90	0.15	16,40,42,49	0
2	6HN	A	301	12/12	0.91	0.10	14,19,21,24	0
2	6HN	B	301	12/12	0.92	0.10	13,19,23,24	0
6	BR	A	308	1/1	0.94	0.29	43,43,43,43	0
4	SIN	B	303	8/8	0.95	0.10	9,13,20,23	0
6	BR	A	307	1/1	0.96	0.25	35,35,35,35	0
4	SIN	A	303	8/8	0.96	0.07	11,12,20,23	0
6	BR	A	306	1/1	0.99	0.16	22,22,22,22	0
3	VVO	B	302	2/2	0.99	0.05	10,10,10,13	0
3	VVO	A	302	2/2	0.99	0.04	10,10,10,13	0
6	BR	B	307	1/1	0.99	0.17	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around 6HN A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around 6HN B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SIN B 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SIN A 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around VVO B 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around VVO A 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.