



wwPDB NMR Structure Validation Summary Report ⓘ

Aug 22, 2026 – 06:23 am BST

PDB ID : 1E08 / pdb_00001e08
Title : Structural model of the [Fe]-Hydrogenase/cytochrome c553 complex combining NMR and soft-docking
Authors : Morelli, X.; Czjzek, M.; Hatchikian, C.E.; Bornet, O.; Fontecilla-Camps, J.C.; Palma, N.P.; Moura, J.J.G.; Guerlesquin, F.
Deposited on : 2000-03-13

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR, THEORETICAL MODEL

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	371	
2	D	88	
3	E	78	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
10	E	HEC	80	1	-
5	A	PDT	4	-	1

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 5058 atoms, of which 882 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called [FE]-HYDROGENASE (LARGE SUBUNIT).

Mol	Chain	Residues	Atoms						Trace
1	A	371	Total	C	H	N	O	S	0
			3387	1783	580	465	527	32	


- Molecule 2 is a protein called [FE]-HYDROGENASE (SMALL SUBUNIT).

Mol	Chain	Residues	Atoms						Trace
2	D	88	Total	C	H	N	O	S	0
			875	457	162	123	132	1	

- Molecule 3 is a protein called CYTOCHROME C553.

Mol	Chain	Residues	Atoms						Trace
3	E	78	Total	C	H	N	O	S	0
			710	354	138	97	114	7	

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		
4	A	1	Total	Fe	S
			8	4	4

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		
4	A	1	Total	Fe	S
			8	4	4
4	A	1	Total	Fe	S
			8	4	4

- Molecule 5 is 1,3-PROPANEDITHIOL (CCD ID: PDT) (formula: C₃H₈S₂).




Mol	Chain	Residues	Atoms		
5	A	1	Total	C	S
			5	3	2

- Molecule 6 is FE (II) ION (CCD ID: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms	
6	A	2	Total	Fe
			2	2

- Molecule 7 is CYANIDE ION (CCD ID: CYN) (formula: CN).



Mol	Chain	Residues	Atoms		
7	A	1	Total	C	N
			2	1	1
7	A	1	Total	C	N
			2	1	1

- Molecule 8 is CARBON MONOXIDE (CCD ID: CMO) (formula: CO).




Mol	Chain	Residues	Atoms		
8	A	1	Total	C	O
			2	1	1
8	A	1	Total	C	O
			2	1	1

- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
9	D	1	Total	Zn
			1	1

- Molecule 10 is HEME C (CCD ID: HEC) (formula: C₃₄H₃₆FeN₄O₄).



Mol	Chain	Residues	Atoms				
10	E	1	Total	C	Fe	N	O
			43	34	1	4	4


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		
11	A	1	Total	H	O
			3	2	1


4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: [FE]-HYDROGENASE (LARGE SUBUNIT)



• Molecule 2: [FE]-HYDROGENASE (SMALL SUBUNIT)



• Molecule 3: CYTOCHROME C553



5 Refinement protocol and experimental data overview

The models were refined using the following method: ?.

Of the 1000 calculated structures, 1 were deposited, based on the following criterion: *LEAST SHIFT VARIATION VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.843
UXNMR	structure solution	UXNMR

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, CYN, CMO, SF4, PDT, HEC, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.47	27/2874 (0.9%)	2.22	114/3885 (2.9%)
2	D	2.23	15/733 (2.0%)	2.75	68/988 (6.9%)
3	E	1.01	4/579 (0.7%)	1.80	21/765 (2.7%)
All	All	1.58	46/4186 (1.1%)	2.27	203/5638 (3.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	7	23
2	D	2	10
3	E	1	3
All	All	10	36

5 of 46 bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	77	GLU	C-N	-44.94	0.94	1.33
1	A	72	CYS	C-N	32.13	1.74	1.33
1	A	27	PHE	C-N	-26.00	1.01	1.33
1	A	65	ALA	C-N	-23.52	1.01	1.33
1	A	65	ALA	N-CA	15.72	1.66	1.46

5 of 203 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	29	GLN	O-C-N	-33.85	77.56	122.59
1	A	28	VAL	O-C-N	25.41	145.84	122.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ILE	O-C-N	23.71	149.38	122.83
2	D	81	GLY	O-C-N	-20.57	110.39	123.35
2	D	120	TYR	O-C-N	20.08	144.41	121.32

5 of 10 chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	54	MET	CA
1	A	60	ILE	CB
1	A	142	CYS	CA
1	A	148	THR	CB
1	A	195	PRO	CA

5 of 33 planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	27	PHE	Peptide,Mainchain
1	A	29	GLN	Mainchain
1	A	41	CYS	Mainchain
1	A	76	CYS	Peptide
1	A	80	ALA	Peptide

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	2807	580	2761	154
11	A	1	2	0	6
2	D	713	162	701	39
3	E	572	138	566	17
4	A	24	0	0	1
5	A	5	0	6	15
6	A	2	0	0	2
7	A	4	0	0	14
8	A	4	0	0	9
All	All	4176	882	4064	184

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 22.

5 of 184 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:72:CYS:C	1:A:73:LEU:N	1.46	1.74
1:A:295:ILE:CD1	1:A:305:ALA:HB2	1.24	1.62
1:A:62:HIS:HD2	2:D:108:LEU:CD1	1.19	1.47
1:A:71:GLN:CD	1:A:385:GLY:HA2	1.19	1.59
1:A:71:GLN:HG2	1:A:385:GLY:CA	1.19	1.66

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/371 (99%)	297 (80%)	52 (14%)	20 (5%)	2	22
2	D	86/88 (98%)	69 (80%)	10 (12%)	7 (8%)	1	13
3	E	76/78 (97%)	68 (89%)	7 (9%)	1 (1%)	12	60
All	All	531/537 (99%)	434 (82%)	69 (13%)	28 (5%)	2	22

5 of 28 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	29	GLN
1	A	36	ILE
1	A	54	MET
1	A	56	GLU
1	A	64	GLU

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/297 (100%)	255 (86%)	42 (14%)	5	44
2	D	76/76 (100%)	66 (87%)	10 (13%)	6	46
3	E	55/55 (100%)	49 (89%)	6 (11%)	8	52
All	All	428/428 (100%)	370 (86%)	58 (14%)	6	45

5 of 58 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	29	GLN
1	A	38	CYS
1	A	42	SER
1	A	47	THR
1	A	60	ILE

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
8	CMO	A	9	6,7	0,1,1	0.00	-
4	SF4	A	1	1	0,12,12	0.00	-
8	CMO	A	10	6,7	0,1,1	0.00	-
10	HEC	E	80	3	50,50,50	0.78	0 (0%)
7	CYN	A	8	8,6,5	0,1,1	0.00	-
5	PDT	A	4	6,7	4,4,4	2.80	2 (50%)
7	CYN	A	7	8,6	0,1,1	0.00	-
4	SF4	A	2	1	0,12,12	0.00	-
4	SF4	A	3	1	0,12,12	0.00	-

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
8	CMO	A	9	6,7	-	-	-
4	SF4	A	1	1	-	-	-
8	CMO	A	10	6,7	-	-	-
10	HEC	E	80	3	68,82,82	1.07	6 (8%)
7	CYN	A	8	8,6,5	-	-	-
5	PDT	A	4	6,7	3,3,3	1.58	0 (0%)
7	CYN	A	7	8,6	-	-	-
4	SF4	A	2	1	-	-	-
4	SF4	A	3	1	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	1	1	-	-	0,6,5,5
10	HEC	E	80	3	1,1,3,7	0,14,54,54	-
5	PDT	A	4	6,7	-	0,2,2,2	-
4	SF4	A	2	1	-	-	0,6,5,5
4	SF4	A	3	1	-	-	0,6,5,5

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	4	PDT	C3-S2	4.77	1.97	1.80
5	A	4	PDT	C1-S1	2.91	1.91	1.80

5 of 6 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	E	80	HEC	C3C-C4C-NC	2.65	113.11	110.15
10	E	80	HEC	CAC-C3C-C2C	2.63	122.66	126.86
10	E	80	HEC	C4C-C3C-C2C	2.29	103.28	106.89
10	E	80	HEC	CAC-C3C-C4C	2.22	128.00	124.92
10	E	80	HEC	C3B-C4B-NB	2.14	112.55	110.17


All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
10	E	80	HEC	NA

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	5
3	E	1
2	D	1

The worst 5 of 7 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	72:CYS	C	73:LEU	N	1.74
1	E	64:TYR	C	65:SER	N	1.19
1	A	202:SER	C	203:PRO	N	1.18
1	A	71:GLN	C	72:CYS	N	1.15
1	A	27:PHE	C	28:VAL	N	1.01

7 Chemical shift validation

No chemical shift data were provided