



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 4, 2026 – 08:47 AM EDT

PDB ID : 4ESV / pdb_00004esv
Title : A New Twist on the Translocation Mechanism of Helicases from the Structure of DnaB with its Substrates
Authors : Itsathitphaisarn, O.; Wing, R.A.; Eliason, W.K.; Wang, J.; Steitz, T.A.
Deposited on : 2012-04-23
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	V	14	36% 57% 7%
2	W	13	15% 69% 15%
3	A	454	2% 55% 27% 9% 8%
3	B	454	2% 52% 30% 11% 5%
3	C	454	56% 31% 8% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	454	
3	E	454	
3	F	454	
3	G	454	
3	H	454	
3	I	454	
3	J	454	
3	K	454	
3	L	454	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GDP	I	501	-	-	X	-
7	ALF	A	504	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 40296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T P*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	V	14	Total	C	N	O	P	0	0	0
			280	140	28	98	14			

- Molecule 2 is a DNA chain called 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T P*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	W	13	Total	C	N	O	P	0	0	0
			260	130	26	91	13			

- Molecule 3 is a protein called Replicative helicase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	418	Total	C	N	O	S	8	1	0
			3255	2036	570	636	13			
3	B	430	Total	C	N	O	S	0	0	0
			3350	2093	591	652	14			
3	C	434	Total	C	N	O	S	0	0	0
			3367	2104	591	658	14			
3	D	431	Total	C	N	O	S	0	0	0
			3333	2084	583	652	14			
3	E	418	Total	C	N	O	S	0	0	0
			3243	2028	565	637	13			
3	F	421	Total	C	N	O	S	0	0	0
			3274	2049	569	643	13			
3	G	421	Total	C	N	O	S	0	0	0
			3259	2037	568	641	13			
3	H	425	Total	C	N	O	S	0	0	0
			3283	2052	572	645	14			
3	I	418	Total	C	N	O	S	0	0	0
			3245	2034	561	637	13			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	J	416	Total	C	N	O	S	0	0	0
			3237	2025	563	636	13			
3	K	421	Total	C	N	O	S	0	0	0
			3267	2044	568	642	13			
3	L	419	Total	C	N	O	S	0	0	0
			3246	2031	565	637	13			

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	W	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	I	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Ca	0	0
			5	5		
5	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Ca	0	0
			1	1		
5	D	2	Total	Ca	0	0
			2	2		
5	E	1	Total	Ca	0	0
			1	1		
5	F	1	Total	Ca	0	0
			1	1		
5	G	1	Total	Ca	0	0
			1	1		
5	I	1	Total	Ca	0	0
			1	1		
5	L	1	Total	Ca	0	0
			1	1		

- Molecule 6 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 28	C 10	N 5	O 11	P 2	0	0
6	C	1	Total 28	C 10	N 5	O 11	P 2	0	0
6	D	1	Total 28	C 10	N 5	O 11	P 2	0	0
6	E	1	Total 28	C 10	N 5	O 11	P 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	F	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	G	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	I	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	J	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	K	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	L	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 7 is TETRAFLUOROALUMINATE ION (CCD ID: ALF) (formula: AlF_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	Al	F	0	0
			5	1	4		
7	C	1	Total	Al	F	0	0
			5	1	4		
7	C	1	Total	Al	F	0	0
			5	1	4		
7	D	1	Total	Al	F	0	0
			5	1	4		
7	E	1	Total	Al	F	0	0
			5	1	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	G	1	Total	Al	F	0	0
			5	1	4		
7	I	1	Total	Al	F	0	0
			5	1	4		
7	J	1	Total	Al	F	0	0
			5	1	4		
7	K	1	Total	Al	F	0	0
			5	1	4		
7	L	1	Total	Al	F	0	0
			5	1	4		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	O	0	0
			1	1		
8	B	1	Total	O	0	0
			1	1		
8	D	1	Total	O	0	0
			1	1		
8	E	1	Total	O	0	0
			1	1		
8	K	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3'

Chain V: 




- Molecule 2: 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3'

Chain W: 




- Molecule 3: Replicative helicase


Chain A: 




- Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase







• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase









4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	149.12Å 180.32Å 279.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.92 – 3.20 47.92 – 3.20	Depositor EDS
% Data completeness (in resolution range)	87.8 (47.92-3.20) 87.7 (47.92-3.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.242 , 0.289 0.239 , 0.281	Depositor DCC
R_{free} test set	6214 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	94.6	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 89.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	40296	wwPDB-VP
Average B, all atoms (Å ²)	140.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, ALF, CA, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	V	0.58	0/307	1.64	4/472 (0.8%)
2	W	0.72	0/285	1.55	3/438 (0.7%)
3	A	0.79	2/3295 (0.1%)	1.10	14/4450 (0.3%)
3	B	0.80	3/3390 (0.1%)	1.19	20/4576 (0.4%)
3	C	0.76	0/3408	1.07	11/4607 (0.2%)
3	D	0.68	0/3372	1.06	13/4557 (0.3%)
3	E	0.68	0/3279	1.04	11/4428 (0.2%)
3	F	0.68	0/3312	1.06	10/4475 (0.2%)
3	G	0.53	0/3296	0.94	7/4454 (0.2%)
3	H	0.58	0/3319	1.00	13/4482 (0.3%)
3	I	0.58	0/3283	0.96	9/4437 (0.2%)
3	J	0.56	0/3273	0.99	15/4420 (0.3%)
3	K	0.62	0/3305	1.01	10/4467 (0.2%)
3	L	0.54	0/3284	0.96	11/4437 (0.2%)
All	All	0.66	5/40408 (0.0%)	1.05	151/54700 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	A	0	4
3	B	0	5
3	C	0	1
3	D	0	2
3	E	0	1
3	F	0	1
3	I	0	1
3	J	0	1
3	K	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	17

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	32	VAL	CA-CB	-5.46	1.51	1.54
3	B	420	ARG	CA-C	5.41	1.60	1.52
3	A	9	ILE	CA-CB	5.33	1.59	1.53
3	B	421	ASN	CB-CG	5.11	1.64	1.52
3	A	169	THR	CA-CB	5.07	1.61	1.53

The worst 5 of 151 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	421	ASN	N-CA-C	-13.28	95.27	112.93
3	B	439	VAL	N-CA-C	10.34	121.30	108.06
3	H	175	MET	N-CA-C	-10.26	101.44	112.93
3	A	168	GLN	N-CA-C	9.80	125.83	109.76
3	K	321	TYR	N-CA-C	9.67	123.21	108.52

There are no chirality outliers.

5 of 17 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	166	LEU	Peptide
3	A	167	VAL	Peptide
3	A	184	THR	Peptide
3	A	320	ASP	Peptide
3	B	151	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	V	280	0	169	11	0
2	W	260	0	157	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3255	0	3296	132	0
3	B	3350	0	3382	185	0
3	C	3367	0	3384	129	0
3	D	3333	0	3337	127	0
3	E	3243	0	3262	115	0
3	F	3274	0	3305	107	0
3	G	3259	0	3274	90	0
3	H	3283	0	3282	114	0
3	I	3245	0	3268	126	0
3	J	3237	0	3265	114	0
3	K	3267	0	3287	119	0
3	L	3246	0	3263	93	0
4	C	12	0	12	0	0
4	E	12	0	12	0	0
4	I	12	0	12	0	0
4	W	12	0	12	4	0
5	A	5	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	2	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	I	1	0	0	0	0
5	L	1	0	0	0	0
6	A	28	0	12	4	0
6	C	28	0	12	4	0
6	D	28	0	12	3	0
6	E	28	0	12	4	0
6	F	28	0	12	4	0
6	G	28	0	12	5	0
6	I	28	0	12	10	0
6	J	28	0	12	6	0
6	K	28	0	12	3	0
6	L	28	0	12	5	0
7	A	5	0	0	2	0
7	C	10	0	0	0	0
7	D	5	0	0	0	0
7	E	5	0	0	0	0
7	G	5	0	0	0	0
7	I	5	0	0	0	0
7	J	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	K	5	0	0	1	0
7	L	5	0	0	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	D	1	0	0	0	0
8	E	1	0	0	0	0
8	K	1	0	0	0	0
All	All	40296	0	40099	1311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 1311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:J:403:ASN:HB3	3:J:404:LYS:CA	1.63	1.28
3:A:145:ARG:HD2	3:B:310:GLN:NE2	1.58	1.18
3:E:14:ILE:HD13	3:E:15:GLU:H	1.10	1.16
2:W:6:DT:H2''	2:W:7:DT:H5'	1.29	1.14
3:J:403:ASN:CB	3:J:404:LYS:HA	1.75	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	413/454 (91%)	369 (89%)	31 (8%)	13 (3%)	3	22
3	B	424/454 (93%)	365 (86%)	38 (9%)	21 (5%)	1	13
3	C	432/454 (95%)	385 (89%)	29 (7%)	18 (4%)	2	16
3	D	425/454 (94%)	376 (88%)	37 (9%)	12 (3%)	4	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	412/454 (91%)	373 (90%)	27 (7%)	12 (3%)	3	24
3	F	415/454 (91%)	377 (91%)	27 (6%)	11 (3%)	4	25
3	G	415/454 (91%)	374 (90%)	37 (9%)	4 (1%)	12	45
3	H	419/454 (92%)	375 (90%)	39 (9%)	5 (1%)	10	42
3	I	412/454 (91%)	375 (91%)	34 (8%)	3 (1%)	18	52
3	J	410/454 (90%)	369 (90%)	37 (9%)	4 (1%)	12	45
3	K	415/454 (91%)	354 (85%)	51 (12%)	10 (2%)	4	28
3	L	413/454 (91%)	372 (90%)	36 (9%)	5 (1%)	10	42
All	All	5005/5448 (92%)	4464 (89%)	423 (8%)	118 (2%)	4	28

5 of 118 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	97	PRO
3	B	154	LYS
3	B	331	SER
3	B	409	LYS
3	B	410	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	353/386 (92%)	290 (82%)	63 (18%)	2	10
3	B	362/386 (94%)	287 (79%)	75 (21%)	1	7
3	C	361/386 (94%)	298 (82%)	63 (18%)	2	10
3	D	356/386 (92%)	298 (84%)	58 (16%)	2	12
3	E	348/386 (90%)	286 (82%)	62 (18%)	2	10
3	F	355/386 (92%)	291 (82%)	64 (18%)	2	9
3	G	351/386 (91%)	302 (86%)	49 (14%)	3	17
3	H	350/386 (91%)	302 (86%)	48 (14%)	3	18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	I	350/386 (91%)	300 (86%)	50 (14%)	3	16
3	J	350/386 (91%)	300 (86%)	50 (14%)	3	16
3	K	352/386 (91%)	295 (84%)	57 (16%)	2	12
3	L	349/386 (90%)	299 (86%)	50 (14%)	3	16
All	All	4237/4632 (92%)	3548 (84%)	689 (16%)	2	12

5 of 689 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	298	VAL
3	J	340	SER
3	H	391	ASP
3	H	293	THR
3	I	316	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 137 such sidechains are listed below:

Mol	Chain	Res	Type
3	K	48	HIS
3	K	225	GLN
3	L	226	ASN
3	E	323	GLN
3	E	259	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 38 ligands modelled in this entry, 14 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GDP	E	501	5,7	29,30,30	1.20	3 (10%)	45,47,47	2.03	10 (22%)
6	GDP	F	501	5,7	29,30,30	1.23	2 (6%)	45,47,47	1.89	11 (24%)
7	ALF	D	503	6	4,4,4	1.30	0	-		
6	GDP	D	501	5,7	29,30,30	1.29	3 (10%)	45,47,47	1.90	7 (15%)
7	ALF	I	503	6	4,4,4	1.29	0	-		
4	MES	E	504	-	12,12,12	1.99	1 (8%)	15,16,16	5.85	8 (53%)
7	ALF	K	502	6	4,4,4	1.45	0	-		
6	GDP	L	501	5,7	29,30,30	1.24	2 (6%)	45,47,47	2.11	12 (26%)
7	ALF	G	502	6	4,4,4	1.45	0	-		
4	MES	C	505	-	12,12,12	2.17	1 (8%)	15,16,16	2.53	7 (46%)
6	GDP	I	501	5,7	29,30,30	1.34	4 (13%)	45,47,47	2.01	17 (37%)
7	ALF	A	504	6	4,4,4	1.27	0	-		
7	ALF	E	503	6	4,4,4	1.30	0	-		
6	GDP	J	501	7	29,30,30	1.19	3 (10%)	45,47,47	1.86	6 (13%)
7	ALF	C	503	6	4,4,4	1.21	0	-		
7	ALF	L	502	6	4,4,4	1.35	0	-		
7	ALF	J	502	6	4,4,4	1.49	0	-		
6	GDP	G	501	7	29,30,30	1.23	2 (6%)	45,47,47	1.86	7 (15%)
4	MES	W	101	-	12,12,12	2.22	1 (8%)	15,16,16	2.29	6 (40%)
6	GDP	K	501	7	29,30,30	1.25	3 (10%)	45,47,47	1.93	9 (20%)
7	ALF	C	504	6	4,4,4	1.25	0	-		
4	MES	I	504	-	12,12,12	2.04	1 (8%)	15,16,16	2.51	6 (40%)
6	GDP	C	501	5,7	29,30,30	1.45	5 (17%)	45,47,47	2.14	12 (26%)
6	GDP	A	502	5,7	29,30,30	1.36	4 (13%)	45,47,47	2.11	11 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GDP	G	501	7	-	3/16/32/32	0/3/3/3
4	MES	C	505	-	-	5/6/14/14	0/1/1/1
6	GDP	E	501	5,7	-	4/16/32/32	0/3/3/3
4	MES	W	101	-	-	2/6/14/14	0/1/1/1
6	GDP	F	501	5,7	-	6/16/32/32	0/3/3/3
6	GDP	K	501	7	-	3/16/32/32	0/3/3/3
6	GDP	D	501	5,7	-	3/16/32/32	0/3/3/3
4	MES	I	504	-	-	3/6/14/14	0/1/1/1
4	MES	E	504	-	-	3/6/14/14	0/1/1/1
6	GDP	C	501	5,7	-	6/16/32/32	0/3/3/3
6	GDP	I	501	5,7	-	3/16/32/32	0/3/3/3
6	GDP	J	501	7	-	4/16/32/32	0/3/3/3
6	GDP	L	501	5,7	-	0/16/32/32	0/3/3/3
6	GDP	A	502	5,7	-	6/16/32/32	0/3/3/3

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	W	101	MES	C8-S	-7.42	1.67	1.77
4	C	505	MES	C8-S	-7.21	1.67	1.77
4	I	504	MES	C8-S	-6.58	1.68	1.77
4	E	504	MES	C8-S	-6.27	1.68	1.77
6	F	501	GDP	C5-C4	3.62	1.48	1.38

The worst 5 of 129 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	504	MES	O2S-S-C8	-12.70	87.53	106.73
4	E	504	MES	O1S-S-C8	-12.24	88.23	106.73
4	E	504	MES	O3S-S-C8	-10.41	85.64	106.00
6	C	501	GDP	C5-C4-N3	-7.50	116.45	128.39
6	A	502	GDP	C5-C4-N3	-7.28	116.81	128.39

There are no chirality outliers.

5 of 51 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	W	101	MES	C8-C7-N4-C5
4	W	101	MES	N4-C7-C8-S

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	C	505	MES	N4-C7-C8-S
4	C	505	MES	C7-C8-S-O1S
4	C	505	MES	C7-C8-S-O3S



There are no ring outliers.



13 monomers are involved in 54 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	501	GDP	4	0
6	F	501	GDP	4	0
6	D	501	GDP	3	0
7	K	502	ALF	1	0
6	L	501	GDP	5	0
6	I	501	GDP	10	0
7	A	504	ALF	2	0
6	J	501	GDP	6	0
6	G	501	GDP	5	0
4	W	101	MES	4	0
6	K	501	GDP	3	0
6	C	501	GDP	4	0
6	A	502	GDP	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	V	14/14 (100%)	-0.52	0	100	100	82, 91, 181, 202	0
2	W	13/13 (100%)	-0.33	0	100	100	92, 107, 167, 208	0
3	A	418/454 (92%)	-0.24	4 (0%)	79	63	29, 100, 162, 214	3 (0%)
3	B	430/454 (94%)	0.00	7 (1%)	70	51	66, 111, 181, 236	0
3	C	434/454 (95%)	-0.34	1 (0%)	91	85	66, 102, 157, 213	0
3	D	431/454 (94%)	-0.24	5 (1%)	76	58	67, 109, 171, 229	0
3	E	418/454 (92%)	-0.31	4 (0%)	79	63	68, 108, 164, 221	0
3	F	421/454 (92%)	-0.33	2 (0%)	87	76	66, 108, 167, 212	0
3	G	421/454 (92%)	0.18	8 (1%)	66	46	100, 165, 252, 322	0
3	H	425/454 (93%)	0.31	19 (4%)	38	24	95, 149, 253, 284	0
3	I	418/454 (92%)	0.28	19 (4%)	38	24	92, 158, 248, 326	0
3	J	416/454 (91%)	0.27	14 (3%)	48	30	59, 177, 285, 314	1 (0%)
3	K	421/454 (92%)	0.12	10 (2%)	59	40	77, 169, 283, 310	0
3	L	419/454 (92%)	0.19	10 (2%)	59	40	112, 174, 246, 301	0
All	All	5099/5475 (93%)	-0.01	103 (2%)	65	45	29, 134, 228, 326	4 (0%)

The worst 5 of 103 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	441	LEU	5.3
3	A	421	ASN	5.0
3	L	386	ILE	4.5
3	J	258	ILE	4.2
3	K	252	LEU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	C	505	12/12	0.64	0.24	232,233,234,235	0
4	MES	W	101	12/12	0.73	0.11	247,250,253,253	0
5	CA	B	501	1/1	0.81	0.17	147,147,147,147	0
5	CA	A	505	1/1	0.82	0.16	117,117,117,117	0
5	CA	A	507	1/1	0.82	0.21	134,134,134,134	0
4	MES	I	504	12/12	0.82	0.12	111,121,151,151	0
7	ALF	G	502	5/5	0.83	0.17	120,127,139,141	0
7	ALF	J	502	5/5	0.85	0.14	119,120,129,130	0
7	ALF	K	502	5/5	0.85	0.10	121,126,137,144	0
6	GDP	L	501	28/28	0.86	0.09	129,162,173,177	0
5	CA	A	506	1/1	0.87	0.13	140,140,140,140	0
6	GDP	K	501	28/28	0.87	0.10	165,178,204,208	0
7	ALF	D	503	5/5	0.89	0.15	80,88,97,109	0
4	MES	E	504	12/12	0.89	0.15	122,127,144,148	0
5	CA	D	504	1/1	0.92	0.08	131,131,131,131	0
5	CA	G	503	1/1	0.92	0.22	141,141,141,141	0
6	GDP	I	501	28/28	0.92	0.09	71,84,103,109	0
6	GDP	J	501	28/28	0.93	0.08	140,157,182,186	0
7	ALF	A	504	5/5	0.93	0.12	84,89,97,97	0
6	GDP	E	501	28/28	0.94	0.09	67,89,110,118	0
6	GDP	A	502	28/28	0.94	0.09	67,75,87,90	0
6	GDP	F	501	28/28	0.95	0.07	71,90,105,111	0
7	ALF	C	504	5/5	0.95	0.12	67,74,92,94	0
5	CA	A	503	1/1	0.95	0.13	103,103,103,103	0
7	ALF	E	503	5/5	0.95	0.11	88,88,100,101	0
5	CA	C	502	1/1	0.95	0.11	101,101,101,101	0
7	ALF	I	503	5/5	0.95	0.08	89,90,95,97	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GDP	C	501	28/28	0.95	0.08	66,69,83,87	0
5	CA	A	501	1/1	0.95	0.19	109,109,109,109	0
6	GDP	G	501	28/28	0.96	0.06	118,157,181,186	0
7	ALF	L	502	5/5	0.96	0.05	118,121,124,126	0
6	GDP	D	501	28/28	0.97	0.06	65,81,90,97	0
7	ALF	C	503	5/5	0.97	0.09	66,70,86,88	0
5	CA	I	502	1/1	0.97	0.10	117,117,117,117	0
5	CA	E	502	1/1	0.97	0.13	110,110,110,110	0
5	CA	D	502	1/1	0.97	0.08	90,90,90,90	0
5	CA	F	502	1/1	0.98	0.11	92,92,92,92	0
5	CA	L	503	1/1	0.98	0.05	155,155,155,155	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around GDP K 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP I 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP J 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP E 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP F 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP G 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.