wwPDB EM Validation Summary Report (i)

Aug 24, 2026 – 05:28 pm BST

PDB ID	:	9FNC / pdb_00009fnc
EMDB ID	:	EMD-50588
Title	:	Half-closed CODH/ACS in the carbonylated state
Authors	:	Ruickoldt, J.; Wendler, P.; Dobbek, H.
Deposited on	:	2024-06-10
Resolution	:	2.21 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	3231 (1.71 - 2.71)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	669	84% 15%
1	B	669	86% 13% •
2	C	730	• 84% 15% •
2	D	730	• 38% • 57%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SF4	A	702	-	-	X	-
5	CMO	C	805	-	-	X	-

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 18850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CO-dehydrogenase.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	669	5130	3254	885	958	33	0	0
1	B	669	5130	3254	885	958	33	0	0

- Molecule 2 is a protein called CO-methylating acetyl-CoA synthase.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	730	5776	3706	969	1072	29	0	0
2	D	311	2466	1599	411	447	9	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	733	ARG	-	expression tag	UNP P83789
C	734	SER	-	expression tag	UNP P83789
D	733	ARG	-	expression tag	UNP P83789
D	734	SER	-	expression tag	UNP P83789

- Molecule 3 is FE(4)-NI(1)-S(4) CLUSTER (CCD ID: XCC) (formula: Fe_4NiS_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	Fe	Ni	S	
3	A	1	9	4	1	4	0
3	B	1	9	4	1	4	0

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
4	A	1	4	2	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
4	A	1	8	4	4	0
4	B	1	4	2	2	0
4	B	1	8	4	4	0
4	C	1	8	4	4	0

- Molecule 5 is CARBON MONOXIDE (CCD ID: CMO) (formula: CO) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
5	A	1	2	1	1	0
5	A	1	2	1	1	0
5	A	1	2	1	1	0
5	A	1	2	1	1	0
5	B	1	2	1	1	0
5	B	1	2	1	1	0
5	B	1	2	1	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
5	B	1	2	1	1	0
5	B	1	2	1	1	0
5	C	1	2	1	1	0
5	C	1	2	1	1	0
5	C	1	2	1	1	0
5	C	1	2	1	1	0
5	C	1	2	1	1	0
5	C	1	2	1	1	0
5	D	1	2	1	1	0
5	D	1	2	1	1	0
5	D	1	2	1	1	0

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
			Total	Ni	
6	C	2	2	2	0

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		AltConf
			Total	Na	
7	C	1	1	1	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
8	A	251	251	251	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
8	B	3	3	3	0
			Total	O	
8	C	3	3	3	0
			Total	O	
8	D	2	2	2	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CO-dehydrogenase

Chain A:

P2 R3 S14 K15 K26 D30 A33 K39 K42 K46 D50 R51 F52 M55 Q60 K64 Q73 R77 G108 H112 E125 E128 G129 K130 D133 R143 T153 I185 D213 I220 G221 N222 A242 D252

P260 E264 I277 H280 L286 K289 C316 N319 E320 M323 R324 H328 V349 V350 D351 V352 Q353 C354 I355 T363 H367 I376 I389 S395 I399 R410 K413 A426 G427 V440 N441 Q442 E443 F464 A465 G466 C467 K471

D475 E476 I479 T480 K483 D489 C497 S498 A499 Q500 A503 F507 L508 L520 K530 P538 S545 C546 V547 D548 N549 T550 R551 D554 V573 A574 S575 A576 P577 E578 K583 I587 W590 L594 P597 V598 G601 L606 V623

Y627 E631 R648 T664 Y670

- Molecule 1: CO-dehydrogenase

Chain B:

P2 S14 W20 R27 A32 E35 K46 F52 R69 D82 P83 S84 K85 K86 S92 L106 H112 R143 K156 E167 S205 I211 Q217 A218 H219 I220 D223 V236 T239 K259 P260 L267 N273 L297

K302 A306 E307 M314 E320 M323 R324 H328 S336 E337 A347 V348 V349 V350 D351 V352 Q353 C354 I355 L359 R369 I389 A406 H416 V425 A426 F464 C467 K471 Q474 L482 M485 L486 D489 T494 F502 P510

L513 L520 I524 E528 E529 L533 Q534 L537 P538 G544 S545 C546 V547 D548 N549 T550 R551 D554 P571 F572 V573 E578 K583 I587 W590 L594 L606 D622 E631 I641 L642 L645 T664 N669 Y670

- Molecule 2: CO-methylating acetyl-CoA synthase

Chain C:

I5 Q9 E16 K19 Y38 L42 I68 E75 V76 R77 M82 E96 F99 R103 Y111 I115 I116 E117 E127 P133 V144 K150 E159 I163 A166 A172 K173 K174 M185 C189 D190 N214 F215 T216 Q217



- Molecule 2: CO-methylating acetyl-CoA synthase

Chain D:



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	150000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	51.0	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2700	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.600	Depositor
Minimum map value	-0.269	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.087	Depositor
Map size (Å)	368.128, 368.128, 368.128	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.719, 0.719, 0.719	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, SF4, NA, XCC, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.14	0/5233	0.33	0/7083
1	B	0.14	0/5233	0.33	0/7083
2	C	0.12	0/5910	0.30	0/8000
2	D	0.12	0/2525	0.30	0/3425
All	All	0.13	0/18901	0.32	0/25591

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5130	0	5149	66	0
1	B	5130	0	5149	50	0
2	C	5776	0	5754	59	0
2	D	2466	0	2487	18	0
3	A	9	0	0	0	0
3	B	9	0	0	0	0
4	A	12	0	0	2	0
4	B	12	0	0	0	0
4	C	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	8	0	0	1	0
5	B	10	0	0	0	0
5	C	12	0	0	2	0
5	D	6	0	0	0	0
6	C	2	0	0	1	0
7	C	1	0	0	0	0
8	A	251	0	0	9	0
8	B	3	0	0	0	0
8	C	3	0	0	0	0
8	D	2	0	0	0	0
All	All	18850	0	18539	188	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:803:NI:NI	5:C:805:CMO:C	1.51	0.84
1:B:464:PHE:HB2	1:B:494:THR:HG22	1.67	0.77
1:A:573:VAL:HG22	1:A:597:PRO:HG2	1.68	0.76
1:A:320:GLU:OE1	1:A:324:ARG:NH1	2.20	0.73
2:C:319:ILE:HG21	2:C:454:LYS:HE3	1.70	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	667/669 (100%)	645 (97%)	21 (3%)	1 (0%)	48 56
1	B	667/669 (100%)	645 (97%)	21 (3%)	1 (0%)	48 56

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	728/730 (100%)	706 (97%)	20 (3%)	2 (0%)	36	41
2	D	309/730 (42%)	301 (97%)	8 (3%)	0	100	100
All	All	2371/2798 (85%)	2297 (97%)	70 (3%)	4 (0%)	44	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	B	347	ALA
2	C	190	ASP
2	C	661	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	548/548 (100%)	533 (97%)	15 (3%)	39	52
1	B	548/548 (100%)	531 (97%)	17 (3%)	35	46
2	C	612/612 (100%)	592 (97%)	20 (3%)	33	44
2	D	256/612 (42%)	254 (99%)	2 (1%)	73	84
All	All	1964/2320 (85%)	1910 (97%)	54 (3%)	40	52

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	513	LEU
2	C	174	LYS
2	C	620	GLU
1	B	547	VAL
2	C	9	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	566	GLN
2	D	54	HIS
1	B	73	GLN
1	B	121	HIS
1	B	353	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 3 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	CMO	B	705	-	0,1,1	-	-	-		
5	CMO	C	810	-	0,1,1	-	-	-		
5	CMO	D	802	-	0,1,1	-	-	-		
5	CMO	B	704	-	0,1,1	-	-	-		
5	CMO	A	706	-	0,1,1	-	-	-		
5	CMO	A	705	-	0,1,1	-	-	-		
5	CMO	C	808	-	0,1,1	-	-	-		
3	XCC	B	701	1	0,12,12	-	-	-		
5	CMO	D	801	-	0,1,1	-	-	-		
4	SF4	B	703	1	0,12,12	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	CMO	A	704	-	0,1,1	-	-	-		
5	CMO	A	707	-	0,1,1	-	-	-		
5	CMO	C	805	6	0,1,1	-	-	-		
4	SF4	A	703	1	0,12,12	-	-	-		
4	SF4	A	702	4,1	0,4,12	-	-	-		
5	CMO	B	706	-	0,1,1	-	-	-		
5	CMO	B	708	-	0,1,1	-	-	-		
5	CMO	D	803	-	0,1,1	-	-	-		
5	CMO	B	707	-	0,1,1	-	-	-		
3	XCC	A	701	1	0,12,12	-	-	-		
4	SF4	B	702	4,1	0,4,12	-	-	-		
4	SF4	C	802	2	0,12,12	-	-	-		
5	CMO	C	809	-	0,1,1	-	-	-		
5	CMO	C	807	-	0,1,1	-	-	-		
5	CMO	C	806	-	0,1,1	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XCC	A	701	1	-	-	0/4/4/4
4	SF4	B	702	4,1	-	-	0/1/1/5
4	SF4	C	802	2	-	-	0/6/5/5
4	SF4	A	703	1	-	-	0/6/5/5
3	XCC	B	701	1	-	-	0/4/4/4
4	SF4	A	702	4,1	-	-	0/1/1/5
4	SF4	B	703	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	707	CMO	1	0
5	C	805	CMO	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	702	SF4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CMO B 705



Ligand CMO C 810



Ligand CMO D 802



Ligand CMO B 704	
 Bond lengths	Bond angles
 Torsions	 Rings

Ligand CMO A 706	
Bond lengths	Bond angles
Torsions	Rings

Ligand CMO A 705	
 Bond lengths	 Bond angles
 Torsions	Rings

Ligand CMO C 808	
Bond lengths	Bond angles
 Torsions	 Rings

Ligand XCC B 701



Ligand CMO D 801



Ligand SF4 B 703



Bond lengths



Bond angles



Torsions



Rings

Ligand CMO A 704	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand CMO A 707	
Bond lengths	Bond angles
 Torsions	 Rings

Ligand CMO C 805	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand SF4 A 703



Bond lengths



Bond angles



Torsions



Rings

Ligand SF4 A 702



Bond lengths



Bond angles



Torsions



Rings

Ligand CMO B 706

Bond lengths

Bond angles



Torsions



Rings

Ligand CMO B 708	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand CMO D 803	
Bond lengths	Bond angles
 Torsions	 Rings

Ligand CMO B 707	
Bond lengths	Bond angles
Torsions	Rings

Ligand XCC A 701



Ligand SF4 B 702



Bond lengths



Bond angles



Torsions



Rings

Ligand SF4 C 802



Ligand CMO C 809	
Bond lengths	Bond angles
 Torsions	 Rings

Ligand CMO C 807	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand CMO C 806	
Bond lengths	Bond angles
 Torsions	 Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-50588. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 256


Y Index: 256


Z Index: 256

6.2.2 Raw map


X Index: 256


Y Index: 256


Z Index: 256

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map


X Index: 261


Y Index: 241


Z Index: 236

6.3.2 Raw map


X Index: 260


Y Index: 242


Z Index: 236

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.087. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_50588_msk_1.map (i)



X



Y



Z

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 108 nm^3; this corresponds to an approximate mass of 98 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.452 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.452 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.21	-	-
Author-provided FSC curve	2.21	2.70	2.29
Unmasked-calculated*	3.12	4.01	3.22

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.12 differs from the reported value 2.21 by more than 10 %

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-50588 and PDB model 9FNC. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.087 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.087).

9.4 Atom inclusion (i)



At the recommended contour level, 92% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.087) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9010	0.6190
A	0.9300	0.6330
B	0.9260	0.6290
C	0.8970	0.6090
D	0.8170	0.5920