wwPDB EM Validation Summary Report (i)

Sep 9, 2026 – 01:55 pm BST

PDB ID	:	6FO0 / pdb_00006fo0
EMDB ID	:	EMD-4286
Title	:	CryoEM structure of bovine cytochrome bc1 in complex with the anti-malarial compound GSK932121
Authors	:	Johnson, R.M.; Amporndanai, K.; O'Neill, P.M.; Fishwick, C.W.G.; Jamson, A.H.; Rawson, S.D.; Hasnain, S.S.; Antonyuk, S.V.; Muench, S.P.
Deposited on	:	2018-02-05
Resolution	:	4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	79% 14% 8%
1	N	480	78% 14% 8%
2	B	453	77% 15% 9%
2	O	453	78% 14% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	379	83% 14%
3	P	379	85% 13%
4	D	325	60% 13% 26%
4	Q	325	62% 11% 26%
5	E	274	15% 58% 7% 35%
5	R	274	15% 58% 7% 35%
6	F	111	82% 6% 12%
6	S	111	82% 6% 12%
7	G	82	78% 12% 10%
7	T	82	80% 10% 10%
8	H	91	5% 67% 29%
8	U	91	8% 69% 29%
9	I	17	24% 71% 29%
9	V	17	18% 71% 29%
10	J	64	5% 84% 5% 11%
10	W	64	6% 86% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	HEC	D	501	X	-	-	-
13	HEC	Q	501	X	-	-	-

2 Entry composition (i)

There are 13 unique types of molecules in this entry. The entry contains 61356 atoms, of which 30362 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome b-c1 complex subunit 1, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	N	443	6753	2142	3325	603	663	20	0	0
1	A	443	6753	2142	3325	603	663	20	0	0

- Molecule 2 is a protein called Cytochrome b-c1 complex subunit 2, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	O	414	6206	1955	3094	549	601	7	0	0
2	B	414	6206	1955	3094	549	601	7	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	305	GLU	GLN	conflict	UNP P23004
B	305	GLU	GLN	conflict	UNP P23004

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	P	370	5933	1973	2997	456	489	18	0	0
3	C	370	5933	1973	2997	456	489	18	0	0

- Molecule 4 is a protein called Cytochrome c1, heme protein, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	Q	239	3751	1216	1848	327	345	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	D	239	3751	1216	1848	327	345	15	0	0

- Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
5	R	178	2658	848	1311	231	260	8	0	0
5	E	178	2658	848	1311	231	260	8	0	0

- Molecule 6 is a protein called Cytochrome b-c1 complex subunit 7.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
6	S	98	1710	547	850	154	157	2	0	0
6	F	98	1710	547	850	154	157	2	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	56	ASP	ASN	conflict	UNP P00129
F	56	ASP	ASN	conflict	UNP P00129

- Molecule 7 is a protein called Cytochrome b-c1 complex subunit 8.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
7	T	74	1255	408	631	117	98	1	0	0
7	G	74	1255	408	631	117	98	1	0	0

- Molecule 8 is a protein called Cytochrome b-c1 complex subunit 6, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
8	U	65	1041	321	512	96	107	5	0	0
8	H	65	1041	321	512	96	107	5	0	0

- Molecule 9 is a protein called Chain I/V.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	V	17	Total	C	H	N	O	0	0
			263	81	136	24	22		
9	I	17	Total	C	H	N	O	0	0
			263	81	136	24	22		

- Molecule 10 is a protein called Cytochrome b-c1 complex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	W	57	Total	C	H	N	O	0	0
			950	312	477	82	79		
10	J	57	Total	C	H	N	O	0	0
			950	312	477	82	79		

- Molecule 11 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					AltConf
11	P	1	Total	C	Fe	N	O	0
			43	34	1	4	4	
11	P	1	Total	C	Fe	N	O	0
			43	34	1	4	4	
11	C	1	Total	C	Fe	N	O	0
			43	34	1	4	4	
11	C	1	Total	C	Fe	N	O	0
			43	34	1	4	4	

- Molecule 12 is 3-chloro-6-(hydroxymethyl)-2-methyl-5-{4-[3-(trifluoromethoxy)phenoxy]phenyl}pyridin-4-ol (CCD ID: G8U) (formula: $C_{20}H_{15}ClF_3NO_4$).



Mol	Chain	Residues	Atoms						AltConf
			Total	C	Cl	F	N	O	
12	P	1	29	20	1	3	1	4	0
12	C	1	29	20	1	3	1	4	0

- Molecule 13 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{36}FeN_4O_4$).



Mol	Chain	Residues	Atoms						AltConf
			Total	C	Fe	N	O		
13	Q	1	43	34	1	4	4		0
13	D	1	43	34	1	4	4		0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome b-c1 complex subunit 1, mitochondrial

Chain N: 78% 14% 8%



• Molecule 1: Cytochrome b-c1 complex subunit 1, mitochondrial

Chain A: 79% 14% 8%



• Molecule 2: Cytochrome b-c1 complex subunit 2, mitochondrial

Chain O: 78% 14% 9%





- Molecule 2: Cytochrome b-c1 complex subunit 2, mitochondrial

Chain B:



- Molecule 3: Cytochrome b

Chain P:



- Molecule 3: Cytochrome b

Chain C:



- Molecule 4: Cytochrome c1, heme protein, mitochondrial

Chain Q:



- Molecule 4: Cytochrome c1, heme protein, mitochondrial

Chain D:



- Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial

Chain R:



- Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial

Chain E:





- Molecule 6: Cytochrome b-c1 complex subunit 7

Chain S:



- Molecule 6: Cytochrome b-c1 complex subunit 7

Chain F:



- Molecule 7: Cytochrome b-c1 complex subunit 8

Chain T:

- Molecule 7: Cytochrome b-c1 complex subunit 8

Chain G:



- Molecule 8: Cytochrome b-c1 complex subunit 6, mitochondrial

Chain U:



- Molecule 8: Cytochrome b-c1 complex subunit 6, mitochondrial

Chain H:

- Molecule 9: Chain I/V

Chain V:



- Molecule 9: Chain I/V

Chain I:



- Molecule 10: Cytochrome b-c1 complex subunit 9

Chain W:



- Molecule 10: Cytochrome b-c1 complex subunit 9

Chain J:



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	232910	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	75	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	79000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.647	Depositor
Minimum map value	-0.279	Depositor
Average map value	0.008	Depositor
Map value standard deviation	0.035	Depositor
Recommended contour level	0.114	Depositor
Map size (Å)	213.00002, 213.00002, 213.00002	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.065, 1.065, 1.065	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, HEM, G8U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.48	0/3500	0.47	0/4752
1	N	0.48	0/3500	0.47	0/4752
2	B	0.48	0/3166	0.48	0/4291
2	O	0.48	0/3166	0.48	0/4291
3	C	0.54	0/3031	0.52	0/4150
3	P	0.54	0/3031	0.52	0/4150
4	D	0.49	0/1962	0.47	0/2665
4	Q	0.49	0/1962	0.47	0/2665
5	E	0.36	0/1373	0.44	0/1861
5	R	0.36	0/1373	0.44	0/1861
6	F	0.49	0/879	0.47	0/1180
6	S	0.49	0/879	0.47	0/1180
7	G	0.47	0/645	0.53	0/873
7	T	0.46	0/645	0.53	0/873
8	H	0.36	0/534	0.43	0/718
8	U	0.36	0/534	0.43	0/718
9	I	0.52	0/129	0.71	0/177
9	V	0.52	0/129	0.71	0/177
10	J	0.45	0/486	0.43	0/655
10	W	0.45	0/486	0.43	0/655
All	All	0.48	0/31410	0.48	0/42644

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
3	P	0	1
5	E	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
5	R	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	15	ASN	Peptide
5	E	174	GLY	Peptide
3	P	15	ASN	Peptide
5	R	174	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	3325	3324	45	0
1	N	3428	3325	3324	48	0
2	B	3112	3094	3090	45	0
2	O	3112	3094	3090	45	0
3	C	2936	2997	2996	37	0
3	P	2936	2997	2996	35	0
4	D	1903	1848	1849	30	0
4	Q	1903	1848	1849	25	0
5	E	1347	1311	1310	12	0
5	R	1347	1311	1310	11	0
6	F	860	850	849	6	0
6	S	860	850	849	5	0
7	G	624	631	630	7	0
7	T	624	631	630	5	0
8	H	529	512	511	3	0
8	U	529	512	511	1	0
9	I	127	136	135	3	0
9	V	127	136	135	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	J	473	477	477	3	0
10	W	473	477	477	3	0
11	C	86	0	60	18	0
11	P	86	0	60	19	0
12	C	29	0	15	3	0
12	P	29	0	15	3	0
13	D	43	0	32	8	0
13	Q	43	0	32	6	0
All	All	30994	30362	30556	344	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 344 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:P:501:HEM:HBA1	11:P:501:HEM:HHA	1.65	0.79
13:D:501:HEC:HBD1	13:D:501:HEC:HHA	1.65	0.78
13:Q:501:HEC:HBD1	13:Q:501:HEC:HHA	1.65	0.78
11:C:501:HEM:HHA	11:C:501:HEM:HBA1	1.66	0.77
1:N:412:SER:OG	10:W:15:ARG:NH2	2.18	0.77

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	441/480 (92%)	398 (90%)	42 (10%)	1 (0%)	43 76
1	N	441/480 (92%)	398 (90%)	42 (10%)	1 (0%)	43 76
2	B	408/453 (90%)	360 (88%)	48 (12%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	O	408/453 (90%)	360 (88%)	48 (12%)	0	100	100
3	C	368/379 (97%)	330 (90%)	38 (10%)	0	100	100
3	P	368/379 (97%)	330 (90%)	38 (10%)	0	100	100
4	D	237/325 (73%)	215 (91%)	22 (9%)	0	100	100
4	Q	237/325 (73%)	216 (91%)	21 (9%)	0	100	100
5	E	170/274 (62%)	151 (89%)	19 (11%)	0	100	100
5	R	170/274 (62%)	151 (89%)	19 (11%)	0	100	100
6	F	96/111 (86%)	91 (95%)	5 (5%)	0	100	100
6	S	96/111 (86%)	90 (94%)	6 (6%)	0	100	100
7	G	72/82 (88%)	64 (89%)	8 (11%)	0	100	100
7	T	72/82 (88%)	64 (89%)	8 (11%)	0	100	100
8	H	63/91 (69%)	58 (92%)	5 (8%)	0	100	100
8	U	63/91 (69%)	58 (92%)	5 (8%)	0	100	100
9	I	15/17 (88%)	8 (53%)	7 (47%)	0	100	100
9	V	15/17 (88%)	8 (53%)	7 (47%)	0	100	100
10	J	55/64 (86%)	53 (96%)	2 (4%)	0	100	100
10	W	55/64 (86%)	53 (96%)	2 (4%)	0	100	100
All	All	3850/4552 (85%)	3456 (90%)	392 (10%)	2 (0%)	49	81

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	N	282	CYS
1	A	282	CYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/394 (93%)	367 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	N	367/394 (93%)	367 (100%)	0	100	100
2	B	325/355 (92%)	325 (100%)	0	100	100
2	O	325/355 (92%)	325 (100%)	0	100	100
3	C	318/327 (97%)	318 (100%)	0	100	100
3	P	318/327 (97%)	318 (100%)	0	100	100
4	D	204/257 (79%)	204 (100%)	0	100	100
4	Q	204/257 (79%)	204 (100%)	0	100	100
5	E	147/228 (64%)	147 (100%)	0	100	100
5	R	147/228 (64%)	147 (100%)	0	100	100
6	F	90/99 (91%)	90 (100%)	0	100	100
6	S	90/99 (91%)	90 (100%)	0	100	100
7	G	66/72 (92%)	66 (100%)	0	100	100
7	T	66/72 (92%)	66 (100%)	0	100	100
8	H	62/85 (73%)	62 (100%)	0	100	100
8	U	62/85 (73%)	62 (100%)	0	100	100
9	I	15/15 (100%)	15 (100%)	0	100	100
9	V	15/15 (100%)	15 (100%)	0	100	100
10	J	48/54 (89%)	48 (100%)	0	100	100
10	W	48/54 (89%)	48 (100%)	0	100	100
All	All	3284/3772 (87%)	3284 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	141	HIS
4	D	121	HIS
1	A	87	ASN
4	D	31	GLN
1	A	85	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
13	HEC	D	501	4	50,50,50	1.75	4 (8%)	68,82,82	1.52	9 (13%)
13	HEC	Q	501	4	50,50,50	1.74	4 (8%)	68,82,82	1.52	9 (13%)
12	G8U	P	503	-	31,31,31	0.93	1 (3%)	41,45,45	1.73	6 (14%)
11	HEM	C	501	-	50,50,50	1.75	7 (14%)	66,82,82	1.47	12 (18%)
11	HEM	P	501	3	50,50,50	1.73	6 (12%)	66,82,82	1.50	13 (19%)
12	G8U	C	503	-	31,31,31	0.94	1 (3%)	41,45,45	1.75	6 (14%)
11	HEM	P	502	3	50,50,50	1.64	6 (12%)	66,82,82	1.37	8 (12%)
11	HEM	C	502	3	50,50,50	1.63	6 (12%)	66,82,82	1.37	8 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	HEC	D	501	4	1/1/3/7	6/14/54/54	-
13	HEC	Q	501	4	1/1/3/7	6/14/54/54	-
12	G8U	P	503	-	-	4/15/15/15	0/3/3/3
11	HEM	C	501	-	-	3/14/54/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	HEM	P	501	3	-	4/14/54/54	-
12	G8U	C	503	-	-	3/15/15/15	0/3/3/3
11	HEM	P	502	3	-	7/14/54/54	-
11	HEM	C	502	3	-	7/14/54/54	-

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	501	HEM	FE-NB	8.23	2.20	1.94
11	P	501	HEM	FE-NB	7.94	2.19	1.94
13	D	501	HEC	C3D-C2D	7.62	1.53	1.36
13	Q	501	HEC	C3D-C2D	7.61	1.52	1.36
11	P	502	HEM	FE-ND	7.41	2.17	1.94

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
13	D	501	HEC	C1D-ND-C4D	6.38	111.67	105.07
13	Q	501	HEC	C1D-ND-C4D	6.34	111.62	105.07
12	C	503	G8U	C27-C20-C19	-6.03	119.55	123.01
12	P	503	G8U	C27-C20-C19	-6.00	119.56	123.01
12	C	503	G8U	C14-C17-C22	-5.64	118.89	123.46

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
13	Q	501	HEC	NA
13	D	501	HEC	NA

5 of 40 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	P	502	HEM	C2D-C3D-CAD-CBD
11	P	502	HEM	C4D-C3D-CAD-CBD
11	C	502	HEM	C2D-C3D-CAD-CBD
11	C	502	HEM	C4D-C3D-CAD-CBD
12	P	503	G8U	F1-C1-O29-C4

There are no ring outliers.

8 monomers are involved in 53 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	D	501	HEC	8	0
13	Q	501	HEC	6	0
12	P	503	G8U	3	0
11	C	501	HEM	10	0
11	P	501	HEM	10	0
12	C	503	G8U	3	0
11	P	502	HEM	9	0
11	C	502	HEM	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEC D 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEC Q 501



Bond lengths



Bond angles



Torsions



Rings

Ligand G8U P 503



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM C 501



Ligand HEM P 501



Ligand G8U C 503



Ligand HEM P 502



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-4286. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map

X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 100


Y Index: 100


Z Index: 100

6.2.2 Raw map

X Index: 100


Y Index: 100


Z Index: 100

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 105



Y Index: 109



Z Index: 58

6.3.2 Raw map

X Index: 93



Y Index: 112



Z Index: 99

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.114. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 214 nm^3; this corresponds to an approximate mass of 194 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	4.07	4.58	4.18
Unmasked-calculated*	4.18	4.76	4.25

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-4286 and PDB model 6FO0. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.114 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.114).

9.4 Atom inclusion (i)



At the recommended contour level, 71% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.114) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7090	0.4070
A	0.7400	0.4170
B	0.7410	0.4210
C	0.7070	0.4110
D	0.7520	0.4170
E	0.5620	0.3480
F	0.7200	0.4130
G	0.7690	0.4120
H	0.6690	0.3570
I	0.5900	0.3800
J	0.6960	0.3980
N	0.7410	0.4160
O	0.7450	0.4200
P	0.7080	0.4110
Q	0.7550	0.4190
R	0.5640	0.3490
S	0.7250	0.4170
T	0.7750	0.4180
U	0.6580	0.3380
V	0.6230	0.3860
W	0.6640	0.3670