Full wwPDB X-ray Structure Validation Report (i)

Aug 4, 2026 – 09:00 PM EDT

PDB ID	:	4HSX / pdb_00004hsx
Title	:	Structure of the L100F mutant of dehaloperoxidase-hemoglobin A from Amphitrite ornata with 4-bromophenol
Authors	:	Thompson, M.K.; Plummer, A.; Franzen, S.
Deposited on	:	2012-10-31
Resolution	:	1.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1985 (1.14-1.10)
Clashscore	190562	2021 (1.14-1.10)
Ramachandran outliers	187476	1978 (1.14-1.10)
Sidechain outliers	187428	1974 (1.14-1.10)
RSRZ outliers	180081	1984 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	85% 13% •
1	B	137	% 85% 14% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BML	A	206	-	X	X	-
5	BML	B	203	-	X	-	-

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 2894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dehaloperoxidase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	137	1156	732	196	220	8	0	11	0
1	B	137	1194	759	196	232	7	0	18	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	PHE	LEU	engineered mutation	UNP Q9NAV8
B	100	PHE	LEU	engineered mutation	UNP Q9NAV8

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0
			Total	O	S		
3	A	1	5	4	1	0	0
			Total	O	S		
3	A	1	5	4	1	0	0
			Total	O	S		
3	B	1	5	4	1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0

- Molecule 5 is 4-BROMOPHENOL (CCD ID: BML) (formula: C_6H_5BrO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	Br	C	O		
5	A	1	8	1	6	1	0	0
			Total	Br	C	O		
5	B	1	8	1	6	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	214	218	218	0	4
			Total	O		
6	B	194	198	198	0	4

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dehaloperoxidase A

Chain A:

- Molecule 1: Dehaloperoxidase A

Chain B:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	58.24Å 67.44Å 69.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.34 – 1.12 48.34 – 1.12	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.34-1.12) 99.9 (48.34-1.12)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	4.62 (at 1.12Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.126 , 0.150 0.126 , 0.150	Depositor DCC
R_{free} test set	5259 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	7.4	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 44.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h,l,k	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	2894	wwPDB-VP
Average B, all atoms (Å^2)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, BML, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.50	12/1202 (1.0%)	1.13	5/1610 (0.3%)
1	B	1.39	8/1268 (0.6%)	1.07	3/1701 (0.2%)
All	All	1.45	20/2470 (0.8%)	1.10	8/3311 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56[A]	THR	N-CA	9.51	1.57	1.46
1	A	56[B]	THR	N-CA	9.51	1.57	1.46
1	B	37	ASN	CA-CB	7.39	1.63	1.53
1	A	56[A]	THR	CA-C	7.25	1.62	1.52
1	A	56[B]	THR	CA-C	7.25	1.62	1.52
1	B	51	LYS	CA-CB	-7.07	1.42	1.53
1	B	114	SER	N-CA	7.01	1.55	1.46
1	B	109	ARG	CZ-NH1	6.61	1.42	1.32
1	A	55[A]	HIS	N-CA	-5.98	1.38	1.46
1	A	55[B]	HIS	N-CA	-5.98	1.38	1.46
1	A	10	ARG	CZ-NH1	5.70	1.40	1.32
1	B	122	ARG	CZ-NH1	5.51	1.40	1.32
1	A	112	GLY	CA-C	5.37	1.58	1.51
1	B	121	ASP	CG-OD1	-5.32	1.15	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	19	ASP	CB-CG	5.15	1.65	1.52
1	A	45	GLU	N-CA	5.11	1.52	1.46
1	A	89	HIS	ND1-CE1	5.08	1.37	1.32
1	A	89	HIS	CG-CD2	5.08	1.41	1.35
1	A	58	LYS	CD-CE	5.03	1.67	1.52
1	B	34	TYR	C-O	5.00	1.30	1.24

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	125	LYS	CB-CG-CD	6.80	126.93	111.30
1	A	10	ARG	NE-CZ-NH2	6.37	124.93	119.20
1	B	122	ARG	CA-CB-CG	6.32	126.74	114.10
1	B	109	ARG	CG-CD-NE	6.10	125.42	112.00
1	A	10	ARG	NH1-CZ-NH2	-5.54	112.11	119.30
1	A	56[A]	THR	N-CA-CB	5.39	117.83	110.01
1	A	56[B]	THR	N-CA-CB	5.39	117.83	110.01
1	B	18	GLN	OE1-CD-NE2	-5.17	117.43	122.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	GLY	Mainchain
1	A	122[A]	ARG	Sidechain
1	A	122[B]	ARG	Sidechain
1	B	65[A]	GLU	Sidechain
1	B	65[B]	GLU	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1156	0	1145	20	0
1	B	1194	0	1189	21	0
2	A	43	0	30	0	0
2	B	43	0	30	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	15	0	0	0	0
3	B	5	0	0	0	0
4	A	6	0	8	1	0
5	A	8	0	2	7	0
5	B	8	0	4	0	0
6	A	218	0	0	4	0
6	B	198	0	0	7	0
All	All	2894	0	2408	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:HG3	6:A:436:HOH:O	1.30	1.25
1:B:87:LYS:HE3	1:B:87:LYS:H	1.08	1.14
1:A:13[B]:LEU:C	1:A:13[B]:LEU:HD13	1.78	1.08
1:A:13[B]:LEU:HD13	1:A:13[B]:LEU:O	1.56	1.02
1:A:100:PHE:CE2	5:A:206:BML:BR4	2.67	1.02
1:A:13[B]:LEU:HD12	1:A:57:GLU:HG3	1.47	0.97
1:A:100:PHE:HE2	5:A:206:BML:BR4	2.05	0.90
1:B:87:LYS:HE3	1:B:87:LYS:N	1.86	0.90
1:A:13[B]:LEU:C	1:A:13[B]:LEU:CD1	2.47	0.85
1:B:55[B]:HIS:CD2	2:B:201:HEM:O2D	2.28	0.85
1:B:87:LYS:H	1:B:87:LYS:CE	1.88	0.84
1:A:13[B]:LEU:CD1	1:A:57:GLU:HG3	2.08	0.83
1:A:37:ASN:HB2	6:A:373:HOH:O	1.77	0.83
1:B:106[A]:GLU:OE2	6:B:452:HOH:O	2.01	0.78
1:B:44[A]:GLN:OE1	6:B:438:HOH:O	2.02	0.76
1:B:55[A]:HIS:HD2	2:B:201:HEM:O2D	1.73	0.72
1:A:21:PHE:CD1	5:A:206:BML:H5	2.26	0.70
1:A:111[B]:SER:OG	1:A:113[B]:GLN:HG3	1.97	0.65
1:A:98:GLU:CD	6:A:424:HOH:O	2.42	0.62
1:A:21:PHE:CE1	5:A:206:BML:C5	2.83	0.62
1:B:20:ILE:HD13	1:B:104[B]:LEU:HD13	1.81	0.62
1:A:37:ASN:HB2	6:A:409:HOH:O	1.99	0.61
1:B:55[B]:HIS:CG	6:B:429:HOH:O	2.57	0.58
1:A:21:PHE:CD1	5:A:206:BML:C5	2.90	0.54
1:B:55[B]:HIS:HB3	6:B:429:HOH:O	2.07	0.53
1:B:52:PHE:O	1:B:56[A]:THR:HG23	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:PHE:CZ	5:A:206:BML:BR4	3.20	0.48
1:B:17:ALA:HB1	1:B:56[B]:THR:HG22	1.97	0.46
1:A:35:PHE:HE2	5:A:206:BML:C2	2.29	0.46
1:B:55[A]:HIS:CD2	2:B:201:HEM:O2D	2.62	0.46
1:B:106[A]:GLU:HG3	6:B:379:HOH:O	2.16	0.46
1:B:55[B]:HIS:CB	6:B:429:HOH:O	2.63	0.45
1:B:63:MET:HE1	1:B:100:PHE:CZ	2.51	0.45
1:A:113[A]:GLN:HB3	4:A:205:GOL:O3	2.16	0.45
1:A:63[B]:MET:HB2	1:A:63[B]:MET:HE2	1.71	0.45
1:B:51:LYS:HD2	1:B:51:LYS:HA	1.85	0.45
1:B:55[B]:HIS:ND1	6:B:429:HOH:O	2.34	0.45
1:B:55[A]:HIS:HD2	2:B:201:HEM:CGD	2.30	0.43
1:A:55[A]:HIS:CE1	1:A:59:VAL:HG21	2.55	0.42
1:B:55[B]:HIS:NE2	2:B:201:HEM:O2D	2.52	0.41
1:B:20:ILE:HD13	1:B:104[B]:LEU:CD1	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/137 (107%)	141 (97%)	5 (3%)	0	100	100
1	B	154/137 (112%)	153 (99%)	1 (1%)	0	100	100
All	All	300/274 (110%)	294 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/117 (109%)	127 (99%)	1 (1%)	73	41
1	B	136/117 (116%)	135 (99%)	1 (1%)	76	46
All	All	264/234 (113%)	262 (99%)	2 (1%)	70	41

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	LYS
1	B	87	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	81	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	SO4	B	202	-	4,4,4	0.79	0	6,6,6	1.35	1 (16%)
3	SO4	A	204	-	4,4,4	0.47	0	6,6,6	0.73	0
2	HEM	A	201	1,6	50,50,50	1.56	10 (20%)	67,82,82	1.22	7 (10%)
2	HEM	B	201	1,6	50,50,50	1.75	13 (26%)	67,82,82	1.27	6 (8%)
3	SO4	A	202	-	4,4,4	0.58	0	6,6,6	0.54	0
5	BML	A	206	1	8,8,8	5.33	4 (50%)	10,10,10	5.90	7 (70%)
3	SO4	A	203	-	4,4,4	0.40	0	6,6,6	0.26	0
4	GOL	A	205	-	5,5,5	0.79	0	5,5,5	0.66	0
5	BML	B	203	1	8,8,8	3.94	3 (37%)	10,10,10	4.01	6 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	201	1,6	-	3/14/54/54	-
2	HEM	B	201	1,6	-	2/14/54/54	-
5	BML	A	206	1	-	-	0/1/1/1
4	GOL	A	205	-	-	0/4/4/4	-
5	BML	B	203	1	-	-	0/1/1/1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	206	BML	BR4-C4	-11.21	1.68	1.90
5	B	203	BML	C3-C4	-9.24	1.20	1.38
5	A	206	BML	C5-C4	7.00	1.52	1.38
5	A	206	BML	C6-C5	5.96	1.48	1.38
5	B	203	BML	BR4-C4	-5.45	1.79	1.90
2	B	201	HEM	FE-NB	3.85	2.06	1.94
2	B	201	HEM	O2A-CGA	-3.77	1.18	1.30
2	A	201	HEM	CBD-CGD	-3.72	1.42	1.50
5	A	206	BML	C3-C4	3.70	1.45	1.38
2	B	201	HEM	O2D-CGD	-3.68	1.18	1.30
2	A	201	HEM	FE-NB	3.65	2.06	1.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	HEM	FE-ND	3.64	2.06	1.94
2	B	201	HEM	C1B-NB	-3.36	1.34	1.40
2	A	201	HEM	C4D-ND	-3.28	1.34	1.40
2	B	201	HEM	CHD-C4C	3.17	1.44	1.38
2	A	201	HEM	FE-NC	3.10	2.05	1.95
2	B	201	HEM	C3B-C4B	-2.86	1.39	1.44
2	A	201	HEM	FE-NA	2.84	2.04	1.95
2	B	201	HEM	FE-NA	2.78	2.04	1.95
5	B	203	BML	C5-C4	2.73	1.43	1.38
2	A	201	HEM	C1B-NB	-2.72	1.35	1.40
2	B	201	HEM	O1D-CGD	2.59	1.30	1.22
2	A	201	HEM	O2A-CGA	-2.57	1.22	1.30
2	B	201	HEM	FE-NC	2.56	2.03	1.95
2	B	201	HEM	CBD-CGD	-2.49	1.44	1.50
2	A	201	HEM	CMB-C2B	-2.36	1.45	1.50
2	B	201	HEM	C4D-C3D	2.21	1.48	1.45
2	A	201	HEM	CHA-C1A	2.12	1.44	1.39
2	A	201	HEM	FE-ND	2.09	2.01	1.94
2	B	201	HEM	CHB-C4A	2.07	1.44	1.39

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	206	BML	C5-C4-C3	-10.76	105.01	121.33
5	A	206	BML	BR4-C4-C3	9.24	131.88	119.30
5	B	203	BML	C2-C3-C4	8.70	129.59	119.18
5	A	206	BML	C6-C5-C4	7.91	128.65	119.18
5	A	206	BML	C2-C3-C4	6.88	127.42	119.18
5	B	203	BML	BR4-C4-C3	6.77	128.52	119.30
2	B	201	HEM	C3B-C2B-C1B	-4.67	102.91	106.41
2	A	201	HEM	C3B-C2B-C1B	-4.03	103.39	106.41
5	A	206	BML	C3-C2-C1	3.83	123.93	119.88
5	B	203	BML	C5-C4-C3	-3.57	115.91	121.33
5	A	206	BML	C5-C6-C1	-3.30	116.39	119.88
2	A	201	HEM	CAD-CBD-CGD	-3.30	104.91	113.67
5	B	203	BML	C3-C2-C1	-3.25	116.44	119.88
2	A	201	HEM	C2B-C1B-NB	2.92	113.20	109.84
5	A	206	BML	BR4-C4-C5	2.78	123.08	119.30
5	B	203	BML	BR4-C4-C5	-2.75	115.55	119.30
2	B	201	HEM	C2B-C1B-NB	2.73	112.97	109.84
2	B	201	HEM	CHA-C4D-ND	2.54	127.50	124.37
3	B	202	SO4	O4-S-O1	2.44	122.30	109.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	201	HEM	C3D-C4D-ND	2.39	112.79	110.17
2	A	201	HEM	CMB-C2B-C1B	2.36	128.72	125.03
5	B	203	BML	C5-C6-C1	-2.34	117.41	119.88
2	A	201	HEM	CMD-C2D-C1D	2.26	128.56	125.03
2	B	201	HEM	O2A-CGA-CBA	2.25	121.12	114.00
2	A	201	HEM	CMC-C2C-C1C	2.16	128.53	124.73
2	B	201	HEM	CAC-C3C-C4C	-2.10	119.82	124.82
2	B	201	HEM	O2A-CGA-O1A	-2.00	118.19	123.33

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	HEM	C3D-CAD-CBD-CGD
2	A	201	HEM	C4D-C3D-CAD-CBD
2	A	201	HEM	C2D-C3D-CAD-CBD
2	B	201	HEM	C4D-C3D-CAD-CBD
2	B	201	HEM	C2D-C3D-CAD-CBD

There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	201	HEM	5	0
5	A	206	BML	7	0
4	A	205	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEM A 201



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM B 201



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	137/137 (100%)	-0.35	0	100	100	3, 8, 16, 29	11 (8%)
1	B	137/137 (100%)	-0.40	1 (0%)	84	88	3, 7, 15, 24	19 (13%)
All	All	274/274 (100%)	-0.37	1 (0%)	88	91	3, 8, 16, 29	30 (10%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	112	GLY	3.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	SO4	A	204	5/5	0.94	0.10	15,15,24,26	5
5	BML	A	206	8/8	0.95	0.09	6,8,9,12	8
5	BML	B	203	8/8	0.95	0.11	7,7,10,12	8
3	SO4	B	202	5/5	0.96	0.08	12,13,17,19	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	A	205	6/6	0.97	0.07	8,11,13,17	0
2	HEM	B	201	43/43	0.99	0.06	7,9,19,27	2
2	HEM	A	201	43/43	0.99	0.05	6,8,19,23	3
3	SO4	A	203	5/5	1.00	0.03	7,8,8,9	0
3	SO4	A	202	5/5	1.00	0.02	5,5,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 201:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around HEM A 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.