



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 4, 2026 – 05:24 PM EDT

PDB ID : 1MRO / pdb_00001mro
Title : METHYL-COENZYME M REDUCTASE
Authors : Ermler, U.; Grabarse, W.
Deposited on : 1997-10-01
Resolution : 1.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1272 (1.18-1.14)
Ramachandran outliers	187476	1246 (1.18-1.14)
Sidechain outliers	187428	1246 (1.18-1.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	548	89% 10% .
1	D	548	89% 11%
2	B	442	91% 9%
2	E	442	93% 7%
3	C	247	90% 9% .
3	F	247	92% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	GOL	A	804	-	-	X	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 21217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METHYL-COENZYME M REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	548	Total	C	N	O	S	19	15	0
			4293	2717	717	839	20			
1	D	548	Total	C	N	O	S	16	13	0
			4290	2718	716	836	20			

- Molecule 2 is a protein called METHYL-COENZYME M REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	442	Total	C	N	O	S	17	13	0
			3352	2123	553	655	21			
2	E	442	Total	C	N	O	S	10	7	0
			3328	2104	552	652	20			


- Molecule 3 is a protein called METHYL-COENZYME M REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	247	Total	C	N	O	S	36	6	0
			2021	1253	358	399	11			
3	F	247	Total	C	N	O	S	34	3	0
			2013	1247	357	398	11			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is FACTOR 430 (CCD ID: F43) (formula: C₄₂H₅₁N₆NiO₁₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	Ni	O	0	0
			62	42	6	1	13		
5	D	1	Total	C	N	Ni	O	0	0
			62	42	6	1	13		

- Molecule 6 is Coenzyme B (CCD ID: TP7) (formula: C₁₁H₂₂NO₇PS).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	S	0
			21	11	1	7	1	1	
6	A	1	Total	C	N	O	P	S	0
			21	11	1	7	1	1	

- Molecule 7 is 1-THIOETHANESULFONIC ACID (CCD ID: COM) (formula: $C_2H_6O_3S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	O	S	0	0
			7	2	3	2		
7	D	1	Total	C	O	S	0	0
			7	2	3	2		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	D	1	Total	C	O	0	0
			6	3	3		
8	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	1	Total	Na	0	0
			1	1		
9	F	1	Total	Na	0	0
			1	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	378	Total	O	0	0
			378	378		
10	B	306	Total	O	0	0
			306	306		
10	C	197	Total	O	0	0
			197	197		
10	D	367	Total	O	0	0
			367	367		
10	E	277	Total	O	0	0
			277	277		
10	F	188	Total	O	0	0
			188	188		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: METHYL-COENZYME M REDUCTASE




• Molecule 1: METHYL-COENZYME M REDUCTASE



• Molecule 2: METHYL-COENZYME M REDUCTASE





• Molecule 2: METHYL-COENZYME M REDUCTASE






● Molecule 3: METHYL-COENZYME M REDUCTASE



● Molecule 3: METHYL-COENZYME M REDUCTASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.72Å 116.88Å 122.58Å 90.00° 92.02° 90.00°	Depositor
Resolution (Å)	30.00 – 1.16	Depositor
% Data completeness (in resolution range)	92.8 (30.00-1.16)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 0.3	Depositor
R, R_{free}	0.197 , 0.207	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	21217	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GL3, TP7, AGM, GOL, MGN, SMC, F43, NA, ZN, MHS, COM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	2/4426 (0.0%)	0.76	6/6006 (0.1%)
1	D	0.38	0/4412	0.67	1/5988 (0.0%)
2	B	0.35	0/3462	0.66	0/4687
2	E	0.36	0/3413	0.67	0/4622
3	C	0.50	2/2083 (0.1%)	0.72	5/2804 (0.2%)
3	F	0.37	0/2061	0.70	2/2775 (0.1%)
All	All	0.39	4/19857 (0.0%)	0.70	14/26882 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	D	0	1
2	B	0	1
2	E	0	3
3	C	0	2
3	F	0	2
All	All	0	14

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	161	ARG	NE-CZ	-13.40	1.18	1.33
1	A	2	ALA	N-CA	-7.58	1.31	1.46
1	A	2	ALA	CA-C	-6.23	1.39	1.52
3	C	161	ARG	CZ-NH1	5.80	1.40	1.32

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	ALA	CA-C-O	16.38	148.65	120.80
1	A	2	ALA	O-C-N	-16.02	97.37	123.00
3	C	161	ARG	NE-CZ-NH2	10.34	128.51	119.20
3	C	161	ARG	CB-CG-CD	5.85	124.76	111.30
1	A	2	ALA	CB-CA-C	-5.58	102.13	110.50

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2	ALA	Mainchain
1	A	257	MHS	Mainchain
1	A	399	SER	Mainchain
1	A	543[A]	ARG	Sidechain
1	A	543[B]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4293	0	4081	43	0
1	D	4290	0	4088	49	0
2	B	3352	0	3352	41	0
2	E	3328	0	3319	26	0
3	C	2021	0	1958	17	0
3	F	2013	0	1947	10	0
4	A	1	0	0	0	0
5	A	62	0	43	2	0
5	D	62	0	43	2	0
6	A	42	0	38	1	0
7	A	7	0	5	1	0
7	D	7	0	5	1	0
8	A	12	0	16	4	0
8	D	12	0	16	3	0
9	C	1	0	0	0	0
9	F	1	0	0	0	0
10	A	378	0	0	7	2
10	B	306	0	0	11	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	C	197	0	0	6	1
10	D	367	0	0	14	2
10	E	277	0	0	1	0
10	F	188	0	0	0	0
All	All	21217	0	18911	168	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:153[B]:LYS:NZ	3:C:153[B]:LYS:CE	1.68	1.55
1:D:371[B]:LEU:CG	1:D:371[B]:LEU:CD2	1.89	1.50
2:E:236[B]:MET:SD	2:E:236[B]:MET:CG	2.08	1.40
2:E:277[B]:LEU:HD23	2:E:282:ILE:HG13	1.20	1.16
2:B:104[B]:GLU:HG3	10:B:7481:HOH:O	1.45	1.13

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:8708:HOH:O	10:D:8435:HOH:O[2_644]	1.87	0.33
10:A:8554:HOH:O	10:D:8672:HOH:O[2_645]	1.98	0.22
10:B:8702:HOH:O	10:C:8682:HOH:O[2_555]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	556/548 (102%)	532 (96%)	24 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	554/548 (101%)	528 (95%)	26 (5%)	0	100	100
2	B	453/442 (102%)	449 (99%)	4 (1%)	0	100	100
2	E	447/442 (101%)	442 (99%)	5 (1%)	0	100	100
3	C	251/247 (102%)	245 (98%)	6 (2%)	0	100	100
3	F	248/247 (100%)	241 (97%)	7 (3%)	0	100	100
All	All	2509/2474 (101%)	2437 (97%)	72 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/433 (104%)	443 (99%)	5 (1%)	65	30
1	D	446/433 (103%)	443 (99%)	3 (1%)	76	48
2	B	354/341 (104%)	350 (99%)	4 (1%)	65	30
2	E	348/341 (102%)	343 (99%)	5 (1%)	59	21
3	C	221/215 (103%)	217 (98%)	4 (2%)	51	13
3	F	218/215 (101%)	214 (98%)	4 (2%)	51	13
All	All	2035/1978 (103%)	2010 (99%)	25 (1%)	61	25

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	15	GLU
2	E	126	GLU
3	F	208	GLU
2	E	98	ASP
2	E	283	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	477	ASN
2	E	318	GLN
3	F	121	GLN
2	E	115	GLN
1	D	42	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	GL3	A	445	1	2,3,4	0.15	0	1,2,4	0.08	0
1	SMC	D	452	1	5,6,7	0.56	0	3,6,8	0.46	0
1	MHS	A	257	1	10,11,12	3.41	4 (40%)	5,14,16	3.56	3 (60%)
1	MHS	D	257	1	10,11,12	2.55	3 (30%)	5,14,16	1.98	2 (40%)
1	AGM	D	271	1	10,11,12	0.47	0	7,13,15	0.23	0
1	MGN	D	400	1	6,9,10	1.04	1 (16%)	7,12,14	2.17	2 (28%)
1	AGM	A	271	1	10,11,12	0.43	0	7,13,15	0.30	0
1	MGN	A	400	1	6,9,10	1.13	1 (16%)	7,12,14	2.20	2 (28%)
1	SMC	A	452	1	5,6,7	0.50	0	3,6,8	0.53	0
1	GL3	D	445	1	2,3,4	0.18	0	1,2,4	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GL3	A	445	1	-	1/1/1/2	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SMC	D	452	1	-	1/3/5/7	-
1	MHS	A	257	1	-	0/5/6/8	0/1/1/1
1	MHS	D	257	1	-	0/5/6/8	0/1/1/1
1	AGM	D	271	1	-	1/10/11/13	-
1	MGN	D	400	1	-	0/7/9/12	-
1	AGM	A	271	1	-	2/10/11/13	-
1	MGN	A	400	1	-	0/7/9/12	-
1	SMC	A	452	1	-	1/3/5/7	-
1	GL3	D	445	1	-	1/1/1/2	-

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	257	MHS	CM-ND1	8.54	1.63	1.46
1	D	257	MHS	CD2-CG	-6.52	1.28	1.36
1	A	257	MHS	CB-CG	4.69	1.58	1.49
1	D	257	MHS	CD2-NE2	3.62	1.43	1.37
1	D	257	MHS	CB-CG	2.68	1.54	1.49

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	MHS	ND1-CE1-NE2	-5.73	109.53	112.94
1	A	400	MGN	CB2-CA-CB1	-4.85	102.98	110.97
1	D	400	MGN	CB2-CA-CB1	-4.56	103.45	110.97
1	A	257	MHS	CD2-NE2-CE1	3.98	110.45	105.24
1	A	257	MHS	CG-CD2-NE2	-3.78	101.77	109.79

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	452	SMC	CA-CB-SG-CS
1	D	452	SMC	CA-CB-SG-CS
1	A	271	AGM	CE2-CD-NE1-CZ
1	A	445	GL3	S-C-CA-N
1	D	445	GL3	S-C-CA-N

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	445	GL3	1	0
1	A	452	SMC	1	0
1	D	445	GL3	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	COM	A	9800	5	6,6,6	1.51	0	8,8,8	1.15	0
8	GOL	D	805	-	5,5,5	0.75	0	5,5,5	0.92	0
6	TP7	A	9000	-	19,20,20	2.30	7 (36%)	24,26,26	1.36	2 (8%)
6	TP7	A	9500	-	19,20,20	2.37	6 (31%)	24,26,26	1.38	3 (12%)
7	COM	D	9700	5	6,6,6	1.58	2 (33%)	8,8,8	1.16	0
8	GOL	D	806	-	5,5,5	0.72	0	5,5,5	0.92	0
8	GOL	A	804	-	5,5,5	0.73	0	5,5,5	0.92	0
8	GOL	A	807	-	5,5,5	0.75	0	5,5,5	0.92	0
5	F43	A	700	1,7	63,71,71	3.05	21 (33%)	73,118,118	1.84	21 (28%)
5	F43	D	800	1,7	63,71,71	4.46	24 (38%)	73,118,118	3.04	29 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	COM	A	9800	5	-	0/4/4/4	-
8	GOL	D	805	-	-	1/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TP7	A	9000	-	-	2/24/24/24	-
6	TP7	A	9500	-	-	3/24/24/24	-
7	COM	D	9700	5	-	0/4/4/4	-
8	GOL	D	806	-	-	1/4/4/4	-
8	GOL	A	804	-	-	1/4/4/4	-
8	GOL	A	807	-	-	1/4/4/4	-
5	F43	A	700	1,7	-	10/28/185/185	-
5	F43	D	800	1,7	-	9/28/185/185	-

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	800	F43	CHB-C1B	-25.50	1.36	1.53
5	D	800	F43	CHD-C1D	12.51	1.63	1.43
5	A	700	F43	NI-NA	12.11	2.18	1.89
5	A	700	F43	NI-NB	10.81	2.15	1.89
5	D	800	F43	C4D-ND	7.58	1.60	1.49

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	800	F43	C4D-ND-C1D	-10.56	92.64	108.46
5	D	800	F43	C3D-C4D-ND	8.36	115.32	102.34
5	D	800	F43	C5D-C2D-C1D	7.38	120.06	110.43
5	D	800	F43	C2D-C1D-ND	7.14	127.49	110.54
5	D	800	F43	C3B-C4B-CHC	-6.47	109.56	123.33

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	700	F43	C3A-CAA-CBA-CCA
5	D	800	F43	C3A-CAA-CBA-CCA
6	A	9500	TP7	O1-C1-N-CA
8	A	804	GOL	O1-C1-C2-O2
8	A	807	GOL	O1-C1-C2-O2


There are no ring outliers.

7 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	9800	COM	1	0
6	A	9500	TP7	1	0
7	D	9700	COM	1	0
8	D	806	GOL	3	0
8	A	804	GOL	4	0
5	A	700	F43	2	0
5	D	800	F43	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand F43 A 700





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	297[B]:TYR	C	298:GLU	N	1.16

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.