



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 03:53 PM EDT

PDB ID : 1NML / pdb_00001nml
Title : Di-haemic Cytochrome c Peroxidase from Pseudomonas nautica 617, form IN (pH 4.0)
Authors : Dias, J.M.; Bonifacio, C.; Alves, T.; Pereira, A.S.; Bourgeois, D.; Moura, I.; Romao, M.J.
Deposited on : 2003-01-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	4% 79% 16% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	401	X	-	-	-
2	HEC	A	402	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CIT	A	501	-	X	-	-
3	CIT	A	503	-	X	-	-
3	CIT	A	504	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called di-haem cytochrome c peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2417	1527	401	476	13			

- Molecule 2 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{36}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	319	Total	O	0	0
			319	319		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: di-haem cytochrome c peroxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.46Å 114.46Å 90.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.20 20.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-2.20) 99.8 (20.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.21Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.179 , 0.199 0.177 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	42.8	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2874	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2476	0.92	7/3372 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	PRO	N-CA-C	-7.55	99.40	110.80
1	A	259	TYR	N-CA-C	7.25	121.66	110.42
1	A	167	GLU	N-CA-C	7.13	121.28	112.58
1	A	256	THR	N-CA-C	7.03	121.53	112.26
1	A	114	GLU	N-CA-C	-5.61	100.83	109.76
1	A	49	ILE	N-CA-C	5.40	116.25	108.48
1	A	248	SER	N-CA-C	5.36	117.20	109.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2417	0	2328	41	0
2	A	86	0	58	1	0
3	A	52	0	20	3	0
4	A	319	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2874	0	2406	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:GLN:HE21	1:A:110:GLN:H	1.04	0.95
1:A:92:GLN:HE21	1:A:106:LYS:H	1.25	0.79
1:A:110:GLN:H	1:A:110:GLN:NE2	1.82	0.75
1:A:110:GLN:HE21	1:A:110:GLN:N	1.85	0.74
1:A:305:GLU:CD	1:A:305:GLU:H	1.97	0.71
1:A:230:LYS:N	1:A:270:GLU:HG3	2.08	0.68
1:A:106:LYS:HD3	1:A:114:GLU:OE2	1.99	0.61
1:A:37:GLU:OE2	1:A:152:MET:HG3	2.01	0.61
1:A:73:TRP:CZ2	1:A:78:ARG:HD3	2.37	0.59
1:A:1:ASP:O	1:A:5:GLU:HG3	2.03	0.58
1:A:181:LEU:O	1:A:186:LYS:HE3	2.08	0.52
1:A:127:LYS:NZ	4:A:739:HOH:O	2.42	0.52
1:A:52:ASN:HD21	1:A:57:VAL:H	1.58	0.51
1:A:97:ARG:NH1	1:A:242:GLU:OE2	2.40	0.50
1:A:122:VAL:HG11	1:A:152:MET:SD	2.52	0.49
1:A:246:ARG:HD2	3:A:501:CIT:O5	2.13	0.48
1:A:119:PRO:O	1:A:123:VAL:HG23	2.14	0.47
1:A:181:LEU:HD22	1:A:298:THR:HG21	1.98	0.46
1:A:64:GLU:CG	1:A:266:TRP:HH2	2.29	0.46
1:A:47:HIS:ND1	1:A:321:PRO:HG3	2.29	0.46
1:A:69:ILE:O	1:A:78:ARG:NH2	2.46	0.46
1:A:216:LEU:HD11	1:A:233:PHE:HB3	1.98	0.45
1:A:246:ARG:CD	3:A:501:CIT:O5	2.64	0.45
1:A:10:MET:CE	1:A:198:THR:HB	2.46	0.45
1:A:52:ASN:ND2	1:A:57:VAL:H	2.15	0.44
1:A:69:ILE:HG13	2:A:401:HEC:CBC	2.47	0.44
1:A:109:VAL:CG1	1:A:157:GLU:HG2	2.48	0.44
1:A:39:PHE:O	1:A:41:PRO:HD3	2.18	0.43
1:A:12:GLU:HG3	4:A:876:HOH:O	2.17	0.43
1:A:93:PHE:CE1	3:A:501:CIT:H41	2.54	0.43
1:A:230:LYS:N	1:A:270:GLU:CG	2.80	0.42
1:A:89:ASN:HB2	1:A:108:PRO:HG2	2.01	0.42
1:A:73:TRP:CD1	1:A:75:LYS:H	2.37	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:ASN:ND2	1:A:80:SER:H	2.18	0.42
1:A:109:VAL:HG12	1:A:157:GLU:HG2	2.02	0.41
1:A:19:PRO:HD3	1:A:88:PHE:CE2	2.55	0.41
1:A:213:PRO:O	1:A:237:GLU:HA	2.20	0.41
1:A:195:ARG:HG3	1:A:284:LEU:CD2	2.51	0.41
1:A:230:LYS:N	1:A:270:GLU:OE2	2.52	0.41
1:A:93:PHE:CD2	1:A:93:PHE:C	2.98	0.41
1:A:73:TRP:CH2	1:A:78:ARG:HD3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/326 (96%)	299 (96%)	12 (4%)	1 (0%)	36	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	231	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	262/269 (97%)	256 (98%)	6 (2%)	44 59

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	ARG
1	A	106	LYS
1	A	110	GLN
1	A	246	ARG
1	A	275	MET
1	A	305	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	HIS
1	A	52	ASN
1	A	79	ASN
1	A	92	GLN
1	A	104	GLN
1	A	110	GLN
1	A	151	ASN
1	A	279	GLN
1	A	286	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	402	1	50,50,50	1.23	5 (10%)	68,82,82	1.54	13 (19%)
3	CIT	A	503	-	12,12,12	2.33	7 (58%)	17,17,17	2.22	9 (52%)
3	CIT	A	504	-	12,12,12	2.57	6 (50%)	17,17,17	2.68	8 (47%)
3	CIT	A	502	-	12,12,12	2.13	1 (8%)	17,17,17	2.16	8 (47%)
3	CIT	A	501	-	12,12,12	2.46	4 (33%)	17,17,17	2.86	11 (64%)
2	HEC	A	401	1	50,50,50	1.25	7 (14%)	68,82,82	3.27	18 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	402	1	1/1/3/7	0/14/54/54	-
3	CIT	A	503	-	-	8/16/16/16	-
3	CIT	A	504	-	-	9/16/16/16	-
3	CIT	A	502	-	-	3/16/16/16	-
3	CIT	A	501	-	-	5/16/16/16	-
2	HEC	A	401	1	1/1/3/7	6/14/54/54	-

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	CIT	O2-C1	-6.37	1.09	1.30
3	A	504	CIT	O2-C1	-5.97	1.11	1.30
3	A	501	CIT	O2-C1	-5.77	1.11	1.30
3	A	501	CIT	O3-C5	3.66	1.34	1.22
3	A	503	CIT	O1-C1	-3.40	1.11	1.22
3	A	503	CIT	O3-C5	3.33	1.33	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	CIT	O3-C5	3.27	1.32	1.22
3	A	503	CIT	O2-C1	-3.15	1.20	1.30
3	A	503	CIT	O4-C5	-3.15	1.20	1.30
3	A	504	CIT	O4-C5	-3.08	1.20	1.30
3	A	504	CIT	C2-C3	2.77	1.57	1.54
2	A	402	HEC	FE-NA	2.76	2.04	1.95
2	A	402	HEC	CMC-C2C	2.72	1.56	1.50
3	A	501	CIT	O4-C5	-2.66	1.22	1.30
3	A	503	CIT	C2-C3	2.63	1.57	1.54
2	A	401	HEC	CMD-C2D	2.53	1.56	1.50
2	A	402	HEC	O2A-CGA	-2.49	1.22	1.30
2	A	401	HEC	CMC-C2C	2.42	1.55	1.50
2	A	402	HEC	FE-NB	2.41	2.02	1.94
3	A	501	CIT	C2-C3	2.34	1.56	1.54
2	A	401	HEC	CMB-C2B	2.26	1.55	1.50
2	A	402	HEC	C1C-C2C	2.20	1.48	1.43
2	A	401	HEC	O2A-CGA	-2.18	1.23	1.30
2	A	401	HEC	CAD-C3D	2.15	1.56	1.51
3	A	503	CIT	O5-C6	2.11	1.28	1.22
3	A	504	CIT	C2-C1	2.09	1.57	1.50
2	A	401	HEC	O2D-CGD	-2.02	1.24	1.30
2	A	401	HEC	FE-NA	2.01	2.01	1.95
3	A	503	CIT	O7-C3	2.01	1.47	1.43
3	A	504	CIT	O5-C6	2.00	1.28	1.22

All (67) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEC	CAD-CBD-CGD	20.62	168.36	113.67
2	A	401	HEC	CBD-CAD-C3D	-8.74	88.35	112.53
2	A	401	HEC	O1D-CGD-CBD	-8.69	95.53	123.09
3	A	504	CIT	O1-C1-C2	-4.81	109.32	122.95
3	A	504	CIT	C3-C2-C1	4.59	126.48	113.92
3	A	504	CIT	O4-C5-C4	4.47	128.52	114.35
3	A	501	CIT	O1-C1-C2	-4.35	110.64	122.95
3	A	501	CIT	O4-C5-C4	4.11	127.38	114.35
3	A	501	CIT	O7-C3-C4	-4.02	100.20	109.38
2	A	401	HEC	CHA-C4D-ND	-3.99	120.12	124.42
3	A	502	CIT	C3-C2-C1	3.94	124.68	113.92
2	A	402	HEC	CHC-C4B-NB	-3.80	119.66	124.37
3	A	503	CIT	O3-C5-C4	-3.76	112.31	122.95
3	A	504	CIT	O3-C5-C4	-3.71	112.45	122.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	CIT	O3-C5-C4	-3.69	112.50	122.95
3	A	502	CIT	O5-C6-C3	-3.51	115.29	122.09
3	A	501	CIT	C4-C3-C6	3.46	117.68	110.03
3	A	501	CIT	O6-C6-C3	3.38	119.62	113.14
2	A	401	HEC	CAD-C3D-C4D	3.35	130.54	124.70
2	A	402	HEC	CHA-C4D-ND	-3.33	120.84	124.42
3	A	502	CIT	O7-C3-C4	-3.27	101.91	109.38
3	A	504	CIT	O6-C6-C3	3.22	119.32	113.14
3	A	501	CIT	O2-C1-O1	3.21	131.60	123.33
3	A	503	CIT	O4-C5-C4	3.21	124.52	114.35
2	A	401	HEC	CBC-CAC-C3C	3.10	120.81	112.42
3	A	501	CIT	O5-C6-C3	-3.06	116.16	122.09
3	A	501	CIT	C4-C3-C2	3.04	117.10	109.31
3	A	502	CIT	O1-C1-C2	-3.02	114.40	122.95
3	A	503	CIT	C3-C2-C1	2.99	122.09	113.92
2	A	401	HEC	C2A-C1A-NA	-2.98	107.45	110.32
2	A	402	HEC	O1A-CGA-CBA	-2.91	113.85	123.09
2	A	402	HEC	CBC-CAC-C3C	2.85	120.15	112.42
3	A	503	CIT	O4-C5-O3	-2.82	116.08	123.33
2	A	402	HEC	CHD-C1D-ND	-2.81	120.90	124.37
3	A	502	CIT	O7-C3-C6	2.77	112.88	108.96
2	A	401	HEC	O1A-CGA-CBA	-2.75	114.37	123.09
3	A	503	CIT	O2-C1-O1	2.73	130.35	123.33
3	A	501	CIT	C3-C2-C1	2.72	121.36	113.92
2	A	401	HEC	CHA-C4D-C3D	2.67	128.67	124.77
3	A	503	CIT	C3-C4-C5	2.62	121.08	113.92
2	A	401	HEC	CAD-C3D-C2D	-2.61	122.97	127.87
2	A	402	HEC	CHA-C1A-C2A	2.60	128.97	124.86
2	A	402	HEC	CBA-CAA-C2A	-2.56	105.46	112.53
3	A	502	CIT	O6-C6-C3	2.54	118.00	113.14
3	A	504	CIT	C3-C4-C5	2.52	120.81	113.92
3	A	504	CIT	O2-C1-C2	2.51	122.29	114.35
3	A	503	CIT	O5-C6-C3	-2.50	117.25	122.09
2	A	402	HEC	CAD-CBD-CGD	2.47	120.22	113.67
3	A	502	CIT	O2-C1-O1	2.46	129.67	123.33
2	A	401	HEC	CHA-C1A-C2A	2.35	128.57	124.86
2	A	401	HEC	C1B-C2B-C3B	-2.34	104.52	106.98
2	A	402	HEC	C4C-CHD-C1D	2.32	131.69	126.25
2	A	401	HEC	CHC-C4B-NB	-2.28	121.55	124.37
3	A	503	CIT	O2-C1-C2	-2.27	107.15	114.35
2	A	401	HEC	CMC-C2C-C3C	2.27	130.44	125.62
2	A	401	HEC	C2B-C1B-NB	2.26	112.52	109.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	CIT	O6-C6-C3	2.25	117.45	113.14
2	A	401	HEC	CAB-C3B-C4B	-2.24	121.87	124.79
2	A	401	HEC	O2A-CGA-CBA	2.19	120.92	114.00
2	A	402	HEC	CHA-C1A-NA	-2.18	122.08	124.45
3	A	502	CIT	O3-C5-C4	2.17	129.08	122.95
2	A	402	HEC	C1A-CHA-C4D	2.13	130.55	126.02
2	A	402	HEC	O2A-CGA-CBA	2.10	120.64	114.00
3	A	504	CIT	C4-C3-C6	2.06	114.59	110.03
3	A	501	CIT	C3-C4-C5	2.04	119.50	113.92
2	A	401	HEC	CAC-C3C-C2C	2.03	130.10	126.89
2	A	402	HEC	C2B-C1B-NB	2.03	112.25	109.90

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	HEC	NA
2	A	402	HEC	NA

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	CIT	C1-C2-C3-O7
3	A	503	CIT	C1-C2-C3-C6
3	A	503	CIT	O7-C3-C4-C5
3	A	503	CIT	C6-C3-C4-C5
3	A	504	CIT	O7-C3-C6-O6
3	A	503	CIT	C1-C2-C3-C4
3	A	503	CIT	C2-C3-C4-C5
2	A	401	HEC	C4C-C3C-CAC-CBC
3	A	504	CIT	C2-C3-C6-O6
2	A	401	HEC	C2C-C3C-CAC-CBC
3	A	501	CIT	C1-C2-C3-O7
3	A	501	CIT	C1-C2-C3-C6
3	A	504	CIT	C3-C4-C5-O3
3	A	503	CIT	C3-C4-C5-O3
3	A	504	CIT	C2-C3-C6-O5
3	A	504	CIT	C4-C3-C6-O6
3	A	502	CIT	C1-C2-C3-C6
3	A	503	CIT	C4-C3-C6-O6
3	A	504	CIT	C4-C3-C6-O5
3	A	502	CIT	C1-C2-C3-O7
3	A	504	CIT	C3-C4-C5-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	HEC	C4B-C3B-CAB-CBB
2	A	401	HEC	C2B-C3B-CAB-CBB
3	A	501	CIT	C3-C4-C5-O3
3	A	501	CIT	C1-C2-C3-C4
2	A	401	HEC	CAD-CBD-CGD-O1D
3	A	504	CIT	O7-C3-C6-O5
3	A	501	CIT	C3-C4-C5-O4
3	A	504	CIT	O1-C1-C2-C3
3	A	502	CIT	C1-C2-C3-C4
2	A	401	HEC	CAA-CBA-CGA-O2A


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	CIT	3	0
2	A	401	HEC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	316/326 (96%)	0.12	14 (4%)	39 36	29, 43, 62, 91	8 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	325	ILE	12.7
1	A	326	PRO	10.3
1	A	324	MET	8.7
1	A	231	GLY	5.6
1	A	1	ASP	4.8
1	A	230	LYS	4.3
1	A	323	ASP	3.7
1	A	74	GLN	3.2
1	A	322	VAL	3.0
1	A	219	LYS	3.0
1	A	286	ASN	2.1
1	A	305	GLU	2.1
1	A	2	ASN	2.0
1	A	183	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CIT	A	504	13/13	0.83	0.17	77,84,91,91	0
3	CIT	A	502	13/13	0.87	0.14	43,53,57,61	0
3	CIT	A	503	13/13	0.90	0.11	64,66,68,69	0
3	CIT	A	501	13/13	0.97	0.06	39,43,48,48	0
2	HEC	A	401	43/43	0.98	0.07	27,30,38,41	0
2	HEC	A	402	43/43	0.99	0.06	31,36,38,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.