



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 12:56 PM EDT

PDB ID : 3PMQ / pdb_00003pmq
Title : Crystal structure of the outer membrane decaheme cytochrome MtrF
Authors : Clarke, T.A.; Edwards, M.J.; Richardson, D.J.
Deposited on : 2010-11-17
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	669	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	670	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	671	X	-	-	-
2	HEC	A	672	X	-	-	-
2	HEC	A	673	X	-	-	-
2	HEC	A	674	X	-	-	-
2	HEC	A	675	X	-	-	-
2	HEC	A	677	X	-	-	-
2	HEC	A	678	X	-	-	-
2	HEC	A	679	X	-	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Decaheme cytochrome c MtrF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	593	Total	C	N	O	S	0	0	0
			4395	2688	791	883	33			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	640	LYS	-	expression tag	UNP Q8EG32
A	641	GLY	-	expression tag	UNP Q8EG32
A	642	GLU	-	expression tag	UNP Q8EG32
A	643	LEU	-	expression tag	UNP Q8EG32
A	644	LYS	-	expression tag	UNP Q8EG32
A	645	LEU	-	expression tag	UNP Q8EG32
A	646	GLU	-	expression tag	UNP Q8EG32
A	647	GLY	-	expression tag	UNP Q8EG32
A	648	LYS	-	expression tag	UNP Q8EG32
A	649	PRO	-	expression tag	UNP Q8EG32
A	650	ILE	-	expression tag	UNP Q8EG32
A	651	PRO	-	expression tag	UNP Q8EG32
A	652	ASN	-	expression tag	UNP Q8EG32
A	653	PRO	-	expression tag	UNP Q8EG32
A	654	LEU	-	expression tag	UNP Q8EG32
A	655	LEU	-	expression tag	UNP Q8EG32
A	656	GLY	-	expression tag	UNP Q8EG32
A	657	LEU	-	expression tag	UNP Q8EG32
A	658	ASP	-	expression tag	UNP Q8EG32
A	659	SER	-	expression tag	UNP Q8EG32
A	660	THR	-	expression tag	UNP Q8EG32
A	661	ARG	-	expression tag	UNP Q8EG32
A	662	THR	-	expression tag	UNP Q8EG32
A	663	GLY	-	expression tag	UNP Q8EG32
A	664	HIS	-	expression tag	UNP Q8EG32
A	665	HIS	-	expression tag	UNP Q8EG32
A	666	HIS	-	expression tag	UNP Q8EG32

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	667	HIS	-	expression tag	UNP Q8EG32
A	668	HIS	-	expression tag	UNP Q8EG32
A	669	HIS	-	expression tag	UNP Q8EG32

- Molecule 2 is HEME C (CCD ID: HEC) (formula: C₃₄H₃₆FeN₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	


- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Ca 1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Decaheme cytochrome c MtrF



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	256.64Å 256.64Å 256.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.20 40.00 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.0 (40.00-3.20) 97.6 (40.00-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 3.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.305 , 0.321 0.304 , 0.324	Depositor DCC
R_{free} test set	2296 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	89.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 112.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	4826	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/4485	0.81	8/6111 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	SER	N-CA-C	8.34	124.69	112.94
1	A	115	CYS	N-CA-C	6.46	115.57	108.14
1	A	158	ASP	N-CA-C	5.82	118.96	109.06
1	A	69	SER	CB-CA-C	-5.67	109.04	115.79
1	A	86	ALA	N-CA-C	5.59	117.60	108.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4395	0	4133	477	0
2	A	430	0	300	55	0
3	A	1	0	0	0	0
All	All	4826	0	4433	490	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 53.

The worst 5 of 490 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:ILE:HG12	1:A:171:GLN:HB3	1.26	1.17
1:A:116:PRO:HB3	1:A:124:ASN:HA	1.29	1.15
1:A:133:ALA:HB1	1:A:134:THR:HB	1.15	1.15
1:A:55:VAL:HG21	1:A:61:ILE:H	1.11	1.15
1:A:527:HIS:HD1	2:A:677:HEC:HBC1	0.98	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	591/669 (88%)	454 (77%)	109 (18%)	28 (5%)	2 14

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	ILE
1	A	72	GLU
1	A	114	SER
1	A	120	VAL
1	A	262	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	479/542 (88%)	457 (95%)	22 (5%)	24	58

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	481	GLN
1	A	574	ASN
1	A	562	LEU
1	A	602	GLN
1	A	131	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	297	ASN
1	A	502	ASN
1	A	324	ASN
1	A	548	GLN
1	A	104	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	676	1	50,50,50	1.83	7 (14%)	68,82,82	1.23	3 (4%)
2	HEC	A	677	1	50,50,50	1.81	6 (12%)	68,82,82	1.40	7 (10%)
2	HEC	A	672	1	50,50,50	1.83	8 (16%)	68,82,82	1.26	4 (5%)
2	HEC	A	679	1	50,50,50	1.76	5 (10%)	68,82,82	1.33	8 (11%)
2	HEC	A	673	1	50,50,50	1.78	5 (10%)	68,82,82	1.28	3 (4%)
2	HEC	A	674	1	50,50,50	1.82	8 (16%)	68,82,82	1.27	5 (7%)
2	HEC	A	675	1	50,50,50	1.81	7 (14%)	68,82,82	1.35	6 (8%)
2	HEC	A	671	1	50,50,50	1.85	6 (12%)	68,82,82	1.27	5 (7%)
2	HEC	A	670	1	50,50,50	1.84	7 (14%)	68,82,82	1.25	5 (7%)
2	HEC	A	678	1	50,50,50	1.91	7 (14%)	68,82,82	1.27	4 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	676	1	-	14/14/54/54	-
2	HEC	A	677	1	1/1/3/7	11/14/54/54	-
2	HEC	A	672	1	1/1/3/7	5/14/54/54	-
2	HEC	A	679	1	1/1/3/7	5/14/54/54	-
2	HEC	A	673	1	1/1/3/7	6/14/54/54	-
2	HEC	A	674	1	1/1/3/7	9/14/54/54	-
2	HEC	A	675	1	1/1/3/7	7/14/54/54	-
2	HEC	A	671	1	1/1/3/7	7/14/54/54	-
2	HEC	A	670	1	1/1/3/7	5/14/54/54	-
2	HEC	A	678	1	1/1/3/7	7/14/54/54	-

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	671	HEC	C3D-C2D	8.16	1.54	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	674	HEC	C3D-C2D	8.08	1.54	1.36
2	A	672	HEC	C3D-C2D	8.05	1.54	1.36
2	A	670	HEC	C3D-C2D	8.01	1.54	1.36
2	A	676	HEC	C3D-C2D	7.98	1.54	1.36

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	679	HEC	C1D-ND-C4D	5.72	111.98	105.21
2	A	674	HEC	C1D-ND-C4D	5.71	111.97	105.21
2	A	677	HEC	C1D-ND-C4D	5.51	111.73	105.21
2	A	672	HEC	C1D-ND-C4D	5.48	111.69	105.21
2	A	673	HEC	C1D-ND-C4D	5.48	111.69	105.21

5 of 9 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	670	HEC	NA
2	A	671	HEC	NA
2	A	672	HEC	NA
2	A	673	HEC	NA
2	A	674	HEC	NA

5 of 76 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	670	HEC	C2D-C3D-CAD-CBD
2	A	670	HEC	C4D-C3D-CAD-CBD
2	A	671	HEC	C1A-C2A-CAA-CBA
2	A	671	HEC	C3A-C2A-CAA-CBA
2	A	671	HEC	C2D-C3D-CAD-CBD

There are no ring outliers.

9 monomers are involved in 55 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	676	HEC	2	0
2	A	677	HEC	12	0
2	A	672	HEC	5	0
2	A	679	HEC	5	0
2	A	674	HEC	3	0


Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	675	HEC	8	0
2	A	671	HEC	10	0
2	A	670	HEC	3	0
2	A	678	HEC	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEC A 676




Bond lengths




Bond angles





Torsions




Rings









Ligand HEC A 673




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 674




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC A 675




Bond lengths




Bond angles




Torsions




Rings




Ligand HEC A 670




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	593/669 (88%)	1.36	172 (29%) 1 1	46, 99, 206, 272	0

The worst 5 of 172 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	86	ALA	11.2
1	A	84	PHE	10.8
1	A	126	HIS	10.5
1	A	628	THR	8.3
1	A	158	ASP	7.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEC	A	679	43/43	0.85	0.26	106,176,195,203	0
3	CA	A	680	1/1	0.94	0.09	101,101,101,101	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HEC	A	678	43/43	0.96	0.17	114,162,187,202	0
2	HEC	A	671	43/43	0.97	0.12	63,86,140,148	0
2	HEC	A	672	43/43	0.97	0.11	40,69,89,104	0
2	HEC	A	676	43/43	0.97	0.12	39,73,127,151	0
2	HEC	A	677	43/43	0.98	0.12	73,103,115,122	0
2	HEC	A	673	43/43	0.98	0.10	29,61,98,106	0
2	HEC	A	675	43/43	0.98	0.10	53,83,105,143	0
2	HEC	A	670	43/43	0.98	0.10	55,90,118,129	0
2	HEC	A	674	43/43	0.99	0.08	31,52,83,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEC A 679:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 678:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 671:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 672:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 676:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 677:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 673:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 675:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC A 670:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.