



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 4, 2026 – 10:50 AM EDT

PDB ID : 3S3A / pdb_00003s3a
Title : Structure of Thermus thermophilus cytochrome ba3 oxidase 120s after Xe de-pressurization
Authors : Luna, V.M.; Fee, J.A.; Deniz, A.A.; Stout, C.D.
Deposited on : 2011-05-18
Resolution : 4.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1029 (4.62-3.90)
Clashscore	190562	1074 (4.62-3.90)
Ramachandran outliers	187476	1001 (4.62-3.88)
Sidechain outliers	187428	1021 (4.66-3.86)
RSRZ outliers	180081	1026 (4.62-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	
2	B	166	
3	C	33	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	HAS	A	801	X	-	-	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	0	0
			4384	2971	704	693	16			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP Q5SJ79
A	-4	HIS	-	expression tag	UNP Q5SJ79
A	-3	HIS	-	expression tag	UNP Q5SJ79
A	-2	HIS	-	expression tag	UNP Q5SJ79
A	-1	HIS	-	expression tag	UNP Q5SJ79
A	0	HIS	-	expression tag	UNP Q5SJ79
A	1	HIS	-	expression tag	UNP Q5SJ79

- Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	166	Total	C	N	O	S	0	0	0
			1298	844	216	234	4			

- Molecule 3 is a protein called Cytochrome c oxidase polypeptide 2A.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	33	Total	C	N	O	0	0	0
			259	179	39	41			

- Molecule 4 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cu	0	0
			1	1		

- # HEM

- Molecule 6 is HEME-AS (CCD ID: HAS) (formula: $\text{C}_{54}\text{H}_{64}\text{FeN}_4\text{O}_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 65	C 54	Fe 1	N 4	O 6	0	0

- Molecule 7 is XENON (CCD ID: XE) (formula: Xe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	4	Total	Xe	0	0
			4	4		

- Molecule 8 is DINUCLEAR COPPER ION (CCD ID: CUA) (formula: Cu₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cu	0	0
			2	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	110.29Å 110.29Å 170.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.57 – 4.25 92.57 – 4.25	Depositor EDS
% Data completeness (in resolution range)	99.7 (92.57-4.25) 99.7 (92.57-4.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.79 (at 4.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.279 , 0.330 0.272 , 0.322	Depositor DCC
R_{free} test set	361 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å ²)	146.8	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	6056	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CU, CUA, XE, HAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.00	12/4541 (0.3%)	0.98	8/6232 (0.1%)
2	B	1.21	15/1335 (1.1%)	1.14	13/1822 (0.7%)
3	C	0.65	0/265	0.89	1/359 (0.3%)
All	All	1.03	27/6141 (0.4%)	1.02	22/8413 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	GLU	CD-OE1	19.20	1.61	1.25
1	A	9	SER	C-N	17.77	1.56	1.33
1	A	510	GLU	CD-OE1	16.45	1.56	1.25
1	A	140	LYS	CE-NZ	11.39	1.83	1.49
2	B	144	GLU	CA-C	11.37	1.66	1.53

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	SER	CA-C-N	13.33	138.76	120.63
1	A	9	SER	C-N-CA	13.33	138.76	120.63
2	B	144	GLU	N-CA-C	12.56	128.14	110.68
1	A	9	SER	O-C-N	-8.93	108.71	123.00
1	A	327	ARG	NE-CZ-NH1	-8.28	113.22	121.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	214	GLY	Mainchain
1	A	9	SER	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4384	0	4494	102	24
2	B	1298	0	1280	38	24
3	C	259	0	279	13	0
4	A	1	0	0	0	0
5	A	43	0	30	2	0
6	A	65	0	62	8	0
7	A	4	0	0	0	0
8	B	2	0	0	0	0
All	All	6056	0	6145	142	24

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:ILE:CD1	1:A:357:ILE:CG1	1.75	1.63
6:A:801:HAS:C13	6:A:801:HAS:C12	1.75	1.60
2:B:165:VAL:CG1	2:B:165:VAL:CB	1.86	1.54
1:A:140:LYS:CE	1:A:140:LYS:NZ	1.83	1.41
1:A:262:ASP:HB2	1:A:511:VAL:HG11	1.43	1.00

The worst 5 of 24 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:SER:C	2:B:143:GLY:N[4_445]	1.17	1.03
1:A:9:SER:O	2:B:143:GLY:CA[4_445]	1.44	0.76
1:A:9:SER:C	2:B:143:GLY:CA[4_445]	1.45	0.75
1:A:10:ARG:N	2:B:143:GLY:N[4_445]	1.57	0.63
1:A:9:SER:CA	2:B:143:GLY:CA[4_445]	1.59	0.61

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	552/568 (97%)	516 (94%)	34 (6%)	2 (0%)	30	66
2	B	164/166 (99%)	152 (93%)	11 (7%)	1 (1%)	21	58
3	C	31/33 (94%)	30 (97%)	1 (3%)	0	100	100
All	All	747/767 (97%)	698 (93%)	46 (6%)	3 (0%)	30	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	GLY
1	A	330	ARG
2	B	87	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	450/462 (97%)	437 (97%)	13 (3%)	37 58
2	B	136/136 (100%)	124 (91%)	12 (9%)	9 30
3	C	26/26 (100%)	24 (92%)	2 (8%)	12 33
All	All	612/624 (98%)	585 (96%)	27 (4%)	25 47

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	19	LEU
2	B	71	VAL
2	B	159	ASN
2	B	69	GLN
2	B	77	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	5	HIS
2	B	40	HIS
2	B	122	ASN
2	B	69	GLN
1	A	254	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	CUA	B	802	2	0,1,1	-	-	-		
5	HEM	A	800	1	50,50,50	1.82	10 (20%)	67,82,82	1.21	7 (10%)
6	HAS	A	801	1	72,72,72	3.69	32 (44%)	87,109,109	3.93	43 (49%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	A	800	1	-	3/14/54/54	-
6	HAS	A	801	1	1/1/8/18	11/42/82/82	-

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	801	HAS	CMC-C2C	10.90	1.73	1.50
6	A	801	HAS	CHA-C4D	10.12	1.62	1.39
6	A	801	HAS	CHB-C1D	8.86	1.56	1.38
6	A	801	HAS	C2A-C3A	8.62	1.55	1.36
6	A	801	HAS	C3B-C2B	8.39	1.53	1.34

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	801	HAS	CHC-C4B-NB	-14.14	106.88	124.37
6	A	801	HAS	CHD-C4A-NA	-14.08	109.12	124.45
6	A	801	HAS	C3B-C4B-NB	13.34	125.17	109.84
6	A	801	HAS	C4B-C3B-C2B	-10.19	90.30	107.44
6	A	801	HAS	C1C-CHC-C4B	8.17	145.46	126.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	801	HAS	NA

5 of 14 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	800	HEM	C2B-C3B-CAB-CBB
5	A	800	HEM	C4B-C3B-CAB-CBB
6	A	801	HAS	O11-C11-C3B-C2B
6	A	801	HAS	C12-C11-C3B-C2B
6	A	801	HAS	C2C-C3C-CAC-CBC

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	800	HEM	2	0
6	A	801	HAS	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	554/568 (97%)	0.07	25 (4%) 38 33	123, 123, 123, 123	0
2	B	166/166 (100%)	0.05	2 (1%) 76 61	123, 123, 123, 123	0
3	C	33/33 (100%)	0.03	2 (6%) 27 27	123, 123, 123, 123	0
All	All	753/767 (98%)	0.07	29 (3%) 43 35	123, 123, 123, 123	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	492	LEU	4.6
1	A	310	LEU	4.3
3	C	15	LEU	3.9
2	B	119	GLU	3.7
1	A	489	PHE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	XE	A	564	1/1	0.77	0.11	123,123,123,123	1
7	XE	A	563	1/1	0.78	0.14	123,123,123,123	1
7	XE	A	565	1/1	0.82	0.15	123,123,123,123	1
7	XE	A	566	1/1	0.91	0.43	123,123,123,123	1
4	CU	A	803	1/1	0.93	0.06	123,123,123,123	0
6	HAS	A	801	65/65	0.93	0.16	123,123,123,123	0
5	HEM	A	800	43/43	0.96	0.11	123,123,123,123	0
8	CUA	B	802	2/2	0.98	0.03	123,123,123,123	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.