



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 10:15 PM EDT

PDB ID : 2ZZB / pdb_00002zzb
Title : Crystal structure of human thioredoxin reductase I and terpyridine platinum(II)
Authors : Lo, Y.C.; Ko, T.P.; Wang, A.H.J.
Deposited on : 2009-02-09
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	% 34% 46% 13% • •
1	B	513	33% 47% 15% 5%
1	C	513	% 34% 47% 14% • •
1	D	513	% 33% 50% 11% • 6%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin reductase 1, cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	0	0
			3776	2397	645	713	21			
1	B	485	Total	C	N	O	S	0	0	0
			3750	2385	640	706	19			
1	C	493	Total	C	N	O	S	0	0	0
			3800	2414	649	716	21			
1	D	484	Total	C	N	O	S	0	0	0
			3741	2379	638	705	19			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP Q16881
A	-12	ALA	-	expression tag	UNP Q16881
A	-11	HIS	-	expression tag	UNP Q16881
A	-10	HIS	-	expression tag	UNP Q16881
A	-9	HIS	-	expression tag	UNP Q16881
A	-8	HIS	-	expression tag	UNP Q16881
A	-7	HIS	-	expression tag	UNP Q16881
A	-6	HIS	-	expression tag	UNP Q16881
A	-5	VAL	-	expression tag	UNP Q16881
A	-4	ASP	-	expression tag	UNP Q16881
A	-3	ASP	-	expression tag	UNP Q16881
A	-2	ASP	-	expression tag	UNP Q16881
A	-1	ASP	-	expression tag	UNP Q16881
A	498	CYS	U	SEE REMARK 999	UNP Q16881
B	-13	MET	-	expression tag	UNP Q16881
B	-12	ALA	-	expression tag	UNP Q16881
B	-11	HIS	-	expression tag	UNP Q16881
B	-10	HIS	-	expression tag	UNP Q16881
B	-9	HIS	-	expression tag	UNP Q16881
B	-8	HIS	-	expression tag	UNP Q16881
B	-7	HIS	-	expression tag	UNP Q16881

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP Q16881
B	-5	VAL	-	expression tag	UNP Q16881
B	-4	ASP	-	expression tag	UNP Q16881
B	-3	ASP	-	expression tag	UNP Q16881
B	-2	ASP	-	expression tag	UNP Q16881
B	-1	ASP	-	expression tag	UNP Q16881
B	498	CYS	U	SEE REMARK 999	UNP Q16881
C	-13	MET	-	expression tag	UNP Q16881
C	-12	ALA	-	expression tag	UNP Q16881
C	-11	HIS	-	expression tag	UNP Q16881
C	-10	HIS	-	expression tag	UNP Q16881
C	-9	HIS	-	expression tag	UNP Q16881
C	-8	HIS	-	expression tag	UNP Q16881
C	-7	HIS	-	expression tag	UNP Q16881
C	-6	HIS	-	expression tag	UNP Q16881
C	-5	VAL	-	expression tag	UNP Q16881
C	-4	ASP	-	expression tag	UNP Q16881
C	-3	ASP	-	expression tag	UNP Q16881
C	-2	ASP	-	expression tag	UNP Q16881
C	-1	ASP	-	expression tag	UNP Q16881
C	498	CYS	U	SEE REMARK 999	UNP Q16881
D	-13	MET	-	expression tag	UNP Q16881
D	-12	ALA	-	expression tag	UNP Q16881
D	-11	HIS	-	expression tag	UNP Q16881
D	-10	HIS	-	expression tag	UNP Q16881
D	-9	HIS	-	expression tag	UNP Q16881
D	-8	HIS	-	expression tag	UNP Q16881
D	-7	HIS	-	expression tag	UNP Q16881
D	-6	HIS	-	expression tag	UNP Q16881
D	-5	VAL	-	expression tag	UNP Q16881
D	-4	ASP	-	expression tag	UNP Q16881
D	-3	ASP	-	expression tag	UNP Q16881
D	-2	ASP	-	expression tag	UNP Q16881
D	-1	ASP	-	expression tag	UNP Q16881
D	498	CYS	U	SEE REMARK 999	UNP Q16881

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is 2,2':6',2''-TERPYRIDINE PLATINUM(II) Chloride (CCD ID: TPT) (formula: $C_{15}H_{11}ClN_3Pt$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C N Pt 19 15 3 1	0	0
4	D	1	Total C N Pt 19 15 3 1	0	0


- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	19	Total O 19 19	0	0
5	B	17	Total O 17 17	0	0
5	C	13	Total O 13 13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	8	Total	O	0	0
			8	8		





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	120.95Å 135.16Å 346.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.20 30.00 – 3.21	Depositor EDS
% Data completeness (in resolution range)	95.3 (30.00-3.20) 95.3 (30.00-3.21)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 3.18Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.242 , 0.305 0.210 , 0.263	Depositor DCC
R_{free} test set	2246 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	74.2	Xtriage
Anisotropy	0.286	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15394	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TPT, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	7/3850 (0.2%)	1.37	40/5211 (0.8%)
1	B	1.20	11/3824 (0.3%)	1.36	34/5177 (0.7%)
1	C	1.04	3/3875 (0.1%)	1.33	33/5245 (0.6%)
1	D	1.11	8/3815 (0.2%)	1.33	32/5166 (0.6%)
All	All	1.12	29/15364 (0.2%)	1.35	139/20799 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
1	C	0	2
1	D	0	1
All	All	0	10

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	17	ILE	CA-CB	10.43	1.67	1.54
1	B	60	VAL	CA-CB	-7.51	1.45	1.54
1	D	300	ILE	CA-CB	7.34	1.62	1.54
1	A	189	CYS	CB-SG	7.08	2.04	1.81
1	D	196	VAL	CA-CB	-6.15	1.46	1.54

The worst 5 of 139 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	VAL	N-CA-C	9.25	121.06	108.11
1	C	452	PHE	N-CA-C	-9.09	102.58	114.31
1	B	183	LEU	N-CA-C	8.82	120.59	110.97
1	C	413	ILE	CB-CA-C	-8.67	102.64	114.00
1	C	453	ALA	N-CA-C	-8.54	101.91	111.14

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	167	TYR	Sidechain
1	A	176	TYR	Sidechain
1	A	200	TYR	Sidechain
1	A	406	PHE	Sidechain
1	B	43	PHE	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3776	0	3779	352	0
1	B	3750	0	3763	340	0
1	C	3800	0	3809	372	2
1	D	3741	0	3750	355	0
2	A	53	0	31	1	0
2	B	53	0	31	2	0
2	C	53	0	31	3	0
2	D	53	0	31	4	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	B	19	0	11	0	0
4	D	19	0	11	3	0
5	A	19	0	0	0	0
5	B	17	0	0	0	0
5	C	13	0	0	1	0
5	D	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	15394	0	15247	1352	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

The worst 5 of 1352 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:CYS:CB	1:A:189:CYS:SG	2.04	1.42
1:A:84:ARG:HH11	1:A:84:ARG:HG3	1.11	1.16
1:C:163:GLU:HG2	1:C:295:ALA:HA	1.29	1.14
1:C:498:CYS:HB3	4:D:501:TPT:H15	1.27	1.12
1:A:163:GLU:HG2	1:A:295:ALA:HA	1.33	1.11

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:497:CYS:O	1:C:497:CYS:O[3_554]	1.59	0.61
1:C:498:CYS:SG	1:C:498:CYS:SG[3_554]	1.70	0.50

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/513 (95%)	418 (86%)	57 (12%)	13 (3%)	4	25
1	B	483/513 (94%)	409 (85%)	64 (13%)	10 (2%)	5	31
1	C	491/513 (96%)	404 (82%)	67 (14%)	20 (4%)	2	17
1	D	482/513 (94%)	407 (84%)	63 (13%)	12 (2%)	4	27
All	All	1944/2052 (95%)	1638 (84%)	251 (13%)	55 (3%)	4	25

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	ASP
1	A	174	LYS
1	A	379	GLU
1	B	184	PHE
1	C	173	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/422 (95%)	332 (83%)	69 (17%)	2	10
1	B	399/422 (94%)	340 (85%)	59 (15%)	3	15
1	C	404/422 (96%)	339 (84%)	65 (16%)	2	13
1	D	398/422 (94%)	340 (85%)	58 (15%)	3	16
All	All	1602/1688 (95%)	1351 (84%)	251 (16%)	2	13

5 of 251 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	396	GLU
1	D	254	ILE
1	C	101	MET
1	D	216	VAL
1	D	340	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	428	ASN
1	D	234	ASN
1	C	129	ASN
1	D	133	GLN
1	D	323	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	D	502	-	4,4,4	0.38	0	6,6,6	0.34	0
2	FAD	A	500	-	58,58,58	1.39	8 (13%)	85,89,89	1.58	17 (20%)
3	SO4	B	502	-	4,4,4	0.41	0	6,6,6	0.20	0
2	FAD	C	500	-	58,58,58	1.75	8 (13%)	85,89,89	1.64	17 (20%)
3	SO4	A	502	-	4,4,4	0.45	0	6,6,6	0.26	0
4	TPT	B	501	-	19,23,24	0.92	0	15,35,38	1.98	4 (26%)
3	SO4	C	502	-	4,4,4	0.42	0	6,6,6	0.14	0
4	TPT	D	501	1	19,23,24	0.94	0	15,35,38	1.95	4 (26%)
2	FAD	B	500	-	58,58,58	1.58	8 (13%)	85,89,89	1.61	15 (17%)
2	FAD	D	500	-	58,58,58	1.74	9 (15%)	85,89,89	1.57	16 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	2/34/50/50	0/6/6/6
2	FAD	C	500	-	-	2/34/50/50	0/6/6/6
4	TPT	B	501	-	-	-	0/5/5/5
4	TPT	D	501	1	-	-	0/5/5/5
2	FAD	B	500	-	-	2/34/50/50	0/6/6/6
2	FAD	D	500	-	-	3/34/50/50	0/6/6/6

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	500	FAD	P-O3P	7.54	1.67	1.59
2	C	500	FAD	P-O3P	7.45	1.67	1.59
2	B	500	FAD	P-O3P	6.19	1.66	1.59
2	C	500	FAD	PA-O3P	5.44	1.65	1.59
2	A	500	FAD	PA-O3P	4.55	1.64	1.59

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	500	FAD	N3A-C2A-N1A	-5.31	120.54	128.58
2	A	500	FAD	N3A-C2A-N1A	-5.23	120.67	128.58
2	B	500	FAD	N3A-C2A-N1A	-5.01	121.00	128.58
2	D	500	FAD	N3A-C2A-N1A	-4.82	121.29	128.58
2	C	500	FAD	C5A-C4A-N3A	-4.75	120.17	126.72

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	FAD	O4B-C4B-C5B-O5B
2	A	500	FAD	C3B-C4B-C5B-O5B
2	B	500	FAD	O4B-C4B-C5B-O5B
2	B	500	FAD	C3B-C4B-C5B-O5B
2	C	500	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

6 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	FAD	1	0
2	C	500	FAD	3	0


Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	SO4	1	0
4	D	501	TPT	3	0
2	B	500	FAD	2	0
2	D	500	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand TPT B 501




Bond lengths



Bond angles




Torsions




Rings


Ligand TPT D 501




Bond lengths




Bond angles




Torsions



Rings





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/513 (95%)	-0.45	4 (0%) 82 67	15, 50, 81, 128	0
1	B	485/513 (94%)	-0.49	2 (0%) 88 79	18, 45, 76, 107	0
1	C	493/513 (96%)	-0.28	3 (0%) 85 73	21, 58, 85, 137	0
1	D	484/513 (94%)	-0.36	3 (0%) 85 73	23, 53, 83, 104	0
All	All	1952/2052 (95%)	-0.39	12 (0%) 85 73	15, 51, 83, 137	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	499	GLY	5.7
1	B	493	LEU	3.8
1	D	493	LEU	3.4
1	C	498	CYS	2.6
1	C	20	GLY	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	D	502	5/5	0.84	0.10	71,73,74,75	0
3	SO4	B	502	5/5	0.86	0.07	81,81,82,82	0
3	SO4	A	502	5/5	0.89	0.07	72,73,75,75	0
3	SO4	C	502	5/5	0.90	0.07	83,83,83,83	0
2	FAD	C	500	53/53	0.93	0.10	43,58,61,62	0
2	FAD	A	500	53/53	0.94	0.09	42,56,69,70	0
4	TPT	B	501	19/20	0.94	0.19	29,36,38,39	19
4	TPT	D	501	19/20	0.94	0.19	42,46,48,48	19
2	FAD	D	500	53/53	0.95	0.08	33,41,57,58	0
2	FAD	B	500	53/53	0.96	0.07	24,32,44,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD C 500:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TPT B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around TPT D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD D 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.